

Universidade Federal da Paraíba
Universidade Federal de Campina Grande
Programa Associado de Pós-Graduação em Matemática
Doutorado em Matemática

Qualitative properties of positive singular solutions to nonlinear elliptic systems with critical exponent

por

Rayssa Helena Aires de Lima Caju

João Pessoa - PB

Fevereiro/2018

Qualitative properties of positive singular solutions to nonlinear elliptic systems with critical exponent

por

Rayssa Helena Aires de Lima Caju [†]

sob orientação do

Prof. Dr. João Marcos Bezerra do Ó

e coorientação do

Prof. Dr. Fernando Codá dos Santos Cavalcanti Marques

Tese apresentada ao Corpo Docente do Programa Associado de Pós-Graduação em Matemática - UFPB/UFCG, como requisito parcial para obtenção do título de Doutor em Matemática.

João Pessoa - PB

Fevereiro/2018

[†]Este trabalho contou com apoio financeiro da CAPES

Catálogo na publicação
Seção de Catalogação e Classificação

C139q Caju, Rayssa Helena Aires de Lima.
Qualitative properties of positive singular solutions
to nonlinear elliptic systems with critical exponent /
Rayssa Helena Aires de Lima Caju. - João Pessoa, 2018.
90 f. : il.

Orientação: João Marcos Bezerra do Ó.
Coorientação: Fernando Codá dos Santos Cavalcanti
Marques.
Tese (Doutorado) - UFPB/CCEN.


1. Matemática. 2. Equações de Schrödinger. 3.
Comportamento assintótico. I. Ó, João Marcos Bezerra
do. II. Marques, Fernando Codá dos Santos Cavalcanti.
III. Título.

UFPB/CCSA


Universidade Federal da Paraíba
Universidade Federal de Campina Grande
Programa Associado de Pós-Graduação em Matemática
Doutorado em Matemática

Área de Concentração: Análise

Aprovada em: 23 de fevereiro de 2018



Prof. Dr. Allan George de Carvalho Freitas




Prof. Dr. Damião Júnio Gonçalves Araújo

Prof. Dra. Marta Calanchi

Prof. Dr. Pedro Eduardo Ubilla Lopez

Prof. Dr. Fernando Codá dos Santos Cavalcanti Marques

Coorientador


Prof. Dr. João Marcos Bezerra do Ó

Orientador

Tese apresentada ao Corpo Docente do Programa Associado de Pós-Graduação em Matemática - UFPB/UFCG, como requisito parcial para obtenção do título de Doutor em Matemática.

Fevereiro/2018

Resumo

Neste trabalho estudaremos o comportamento assintótico de soluções positivas do seguinte sistema elíptico acoplado de equações de Schrödinger não lineares

$$\Delta_g u_i - \sum_{j=1}^2 A_{ij}(x) u_j + \frac{n(n-2)}{4} |\mathcal{U}|^{\frac{4}{n-2}} u_i = 0$$

definido em $B_1(0) \setminus \{0\}$ para $n \geq 3$, onde g é uma métrica Riemanniana na bola unitária e o potencial A é um mapa de classe C^1 tal que $A_{ij}(x)$ é uma matriz simétrica para cada x pertencente a $B_1(0)$. Do ponto de vista da geometria conforme, o sistema acima é uma extensão natural de equações do tipo Yamabe.

Abordaremos o problema assumindo primeiramente que g é a métrica euclidiana e que o potencial A é identicamente nulo. Nesse caso iremos provar que as soluções do nosso problema são assintóticas ao que chamaremos de soluções do tipo Fowler. No caso geral, iremos demonstrar que o mesmo resultado inserindo algumas restrições sobre o potencial e assumindo que a dimensão é menor ou igual a cinco.

Palavras-chave: Soluções do tipo Fowler; Comportamento assintótico; Sistemas do tipo Yamabe.

Abstract

In this work we study the asymptotic behavior to positive solutions of the following coupled elliptic system of nonlinear Schrödinger equations

$$\Delta_g u_i - \sum_{j=1}^2 A_{ij}(x) u_j + \frac{n(n-2)}{4} |\mathcal{U}|^{\frac{4}{n-2}} u_i = 0$$

which are defined in the punctured unit ball $B_1(0) \setminus \{0\}$ for $n \geq 3$. Here g is a Riemannian metric on the unit ball and the potential A is assumed a C^1 map such that $A_{ij}(x)$ is a symmetrical matrix for each x in $B_1(0)$. From the viewpoint of conformal geometry, this systems are pure extensions of Yamabe-type equations.

We will approach the problem assuming first that g is the euclidian metric and the potential A vanishes. In this case we are able to prove that the solutions of our problem are asymptotics to what we call Fowler-type solutions. In the general case we will prove the same result by putting some restrictions on the potential and assuming that the dimension is less or equal to five.

Keywords: Fowler-type solutions; Asymptotic behavior; Yamabe-type system.

Agradecimentos

A Deus.

Ao Professor João Marcos Bezerra do Ó por ter orientado este trabalho, pelo conhecimento transmitido, por me motivar a enfrentar novos desafios e, principalmente, pela confiança depositada.

A Professora Flávia Jerônimo pela ajuda nos momentos de dificuldade, apoio, incentivo, confiança, e por ter acreditado em mim sempre.

Ao Professor Fernando Codá Marques por ter me recebido em Princeton, pela paciência, pelo encorajamento e por todo o conhecimento transmitido.

Ao Professor Almir Silva Santos pela colaboração e por inúmeras contribuições.

A minha família, em especial a minha avó, pela cobrança, carinho, apoio e ajuda em todas as decisões.

Ao meu companheiro Pedro Gaspar pelo encorajamento, apoio, confiança e paciência durante os momentos mais obscuros dessa jornada.

Aos meus amigos ao longo dessa caminhada na UFPB em especial à Ageu, Diego, Ellen, Eudes, José Carlos, Leon, Rainelly, Ricardo e Victor pela amizade e companheirismo.

A turma brasileira em Princeton.

Aos professores da UFPB, em especial a Antônio Sales e Sergio de Albuquerque, que me receberam tão jovem na universidade e aos professores Alexandre Simas, Everaldo Souto, Jacqueline Rojas, Manassés Xavier, Sergio de Albuquerque e Uberlândio Severo, que acompanharam toda a minha trajetória a partir da graduação.

A banca examinadora: Prof. por aceitarem participar da avaliação deste trabalho.

A CAPES - Coordenação de Aperfeiçoamento de Pessoal de Nível superior, pelo apoio financeiro.

“Sou metade agonia, metade esperança.”

Jane Austen

Dedicatória

A minha avó e ao meu noivo.

Contents

Introduction	1
Notation and terminology	5
1 Fowler solutions and Jacobi fields	7
1.1 The linearized equation	9
2 Classification result for the Limit system	11
2.1 Radial Symmetry	12
2.2 ODE Analysis	14
2.3 Removable singularity result	16
2.4 Classification Result	19
2.5 Jacobi Fields for the Limit System	21
3 Coupled elliptic system in the punctured ball	25
3.1 Upper bound	25
3.2 Pohozaev invariant and lower bound	33
3.3 Simple convergence to a radial solution	38
4 Yamabe-type system in the punctured ball	45
4.1 Upper bound near a singularity	45
4.2 Pohozaev invariants and removable singularities	56
4.3 Convergence to a Radial Solution	70
References	78

List of Figures

1.1	Level sets	8
-----	----------------------	---

Introduction

In this thesis our main purpose is to study the asymptotic behavior for positive solutions to the following coupled elliptic system of nonlinear Schrödinger equations

$$\Delta_g u_i - \sum_{j=1}^2 A_{ij}(x) u_j + \frac{n(n-2)}{4} |\mathcal{U}|^{\frac{4}{n-2}} u_i = 0 \quad (1)$$

which are defined in the punctured ball $\Omega = B_1(0) \setminus \{0\}$, where g is a smooth Riemannian metric on the unit ball $B_1(0) \subset \mathbb{R}^n$, $n \geq 3$ and $|\mathcal{U}|^2 = u_1^2 + u_2^2$. Here A is a C^1 map from $B_1(0)$ to the vector space of symmetrical 2×2 real matrices $M_2^s(\mathbb{R})$, where we will write $A = (A_{ij})_{i,j}$ and each A_{ij} is a C^1 real value function.

The critical system (1) is weakly coupled by the matrix A , and strongly coupled by the Gross-Pitaevskii type nonlinearity in the right-hand side of (1). Systems involving this type of nonlinearities are strongly related to two branches of mathematical physics. They appear in the Hartree-Fock theory for double condensates, a binary mixture of Bose-Einstein condensates in two different hyperfine states and they also arise in the study of incoherent solitons in nonlinear optics. We refer the reader to [1],[2] and [8].

In the last few years there has been considerable interest in this type of systems. O. Druet, E. Hebey and J. Vetois in [7] studied stability properties for systems of the form (1) in a compact Riemannian manifold when the potential A is less, in the sense of bilinear forms, than the geometric threshold potential of the conformal laplacian. Consequently, from the viewpoint of conformal geometry, our systems are pure extensions of Yamabe-type equations in the strongly coupled regime and the consideration of such elliptic systems provides a natural background for the interplay between geometry and asymptotic analysis.

Yamabe-type problems are also widely studied in the singular case, in which one is

interested in finding complete conformal metrics with constant scalar curvature in non-compact manifolds with a simple structure at infinity. Having in mind the connections between geometry and asymptotic analysis it is natural to study properties such as existence and the asymptotic behavior of solutions to the system (1) in the singular case.

Recall that, from the analytic viewpoint, the simplest instance of the singular Yamabe problem consists in finding positive solutions to

$$\Delta_g u - \frac{n-2}{4(n-1)} R_g u + \frac{n(n-2)}{4} u^{\frac{n+2}{n-2}} = 0, \quad (2)$$

in the punctured unit ball endowed with a metric g which blows-up at the origin.

The case of the flat metric was first studied by L. Caffarelli, B. Gidas and J. Spruck in [3] in the punctured unit ball and they proved that, when 0 is a nonremovable singularity, the local models are given by the radial Fowler solutions. Remember that u_0 is a *Fowler solution* if it satisfies

$$\Delta u_0 + \frac{n(n-2)}{4} u_0^{\frac{n+2}{n-2}} = 0 \quad \text{in } \mathbb{R}^n \setminus \{0\}, \quad (3)$$

which blows-up at origin. Their proof relies on a complicated version of the Alexandrov reflection method, and it was later simplified by N. Korevaar et al in [11].

We begin by considering the system (1) in its simplest form, which will be a natural generalization of the singular Yamabe problem in the flat case. Assuming that the matrix A is identically zero and that g is the euclidian metric, the system (1) becomes

$$\Delta u_i + \frac{n(n-2)}{4} |\mathcal{U}|^{\frac{4}{n-2}} u_i = 0 \quad (4)$$

in the punctured ball $B_1^n(0) \setminus \{0\}$.

Firstly we describe what we mean by a positive singular solution to the general system (1), in particular to (4). We say that \mathcal{U} is a *positive* solution if each coordinate u_i is positive and that it is *singular* if the following holds

$$\lim_{|x| \rightarrow 0} |\mathcal{U}|(x) = +\infty. \quad (5)$$

Our main goal is to prove that singular solutions of (4) are asymptotic to what we call *Fowler-type solutions*, that is, solutions of the limit system

$$\Delta u_i + \frac{n(n-2)}{4} |\mathcal{U}|^{\frac{4}{n-2}} u_i = 0 \quad \text{in } \mathbb{R}^n \setminus \{0\}. \quad (6)$$

Our main result for system (4) is the following

Theorem 0.1 *Suppose that \mathcal{U} is a solution of the system (4) in the punctured ball $B_1^n(0) \setminus \{0\}$. Then there exists a Fowler-type solution $\mathcal{U}_0 = u_0 \Lambda$ of (6), where u_0 is a Fowler solution such that*

$$\mathcal{U}(x) = (1 + O(|x|^\alpha)) \mathcal{U}_0(x) \quad (7)$$

as $x \rightarrow 0$, for some $\alpha > 0$.

In light of the analogy between the singular problem (2) and the system (1), we are led to analyze the properties of our local model, the Fowler-type solutions.

It was proved by the authors in [7] the following classification result for entire solutions of the system

$$\Delta u_i + \frac{n(n-2)}{4} |\mathcal{U}|^{\frac{4}{n-2}} u_i = 0 \quad \text{in } \mathbb{R}^n, \quad (8)$$

Theorem A [7] *Let \mathcal{U} be a nonnegative C^2 -solution of (8). Then there exist $x_0 \in \mathbb{R}^n$, $\mu > 0$, and $\Lambda \in \mathbb{S}_+^1$, such that*

$$\mathcal{U}(x) = \left(\frac{2\mu}{1 + \mu^2 |x - x_0|^2} \right)^{\frac{n-2}{2}} \Lambda$$

for all $x \in \mathbb{R}^n$, where \mathbb{S}_+^1 consists of the elements in the unit sphere in \mathbb{R}^p , with nonnegative coordinates.

This result inspired us to ask whether any singular nonnegative solution is of the form $u_0 \Lambda$, where now u_0 is a Fowler solution and Λ is a vector in the unit sphere with nonnegative coordinates. Indeed, we are able to prove the following result

Theorem 0.2 *Suppose that \mathcal{U}_0 is a nonnegative singular solution for (6). Then there exist $\Lambda \in \mathbb{S}_+^1$ and u_0 a Fowler solution such that*

$$\mathcal{U}_0 = u_0 \Lambda.$$

where \mathbb{S}_+^1 consists of the elements in the unit sphere in \mathbb{R}^2 , with nonnegative coordinates.

Classification results for singular solutions for coupled elliptic systems were conjectured by Z. Chen and C-S. Lin in [5] for a similar system. This classification will allow us to study all the properties that we need for solutions of the limit system based in the properties of Fowler solution.

Using this classification result, the strategy to prove that the solutions of (4) are asymptotic to Fowler-type solution is to get an upper and lower bound. The upper bound will follow from a more precise universal bound for weak supersolutions which are smooth in a open set which is inspired by the works of [18], [16] and [12]. The lower bound depends in a more delicate way of the solution. This finishes the study of the solutions of (4).

While Korevaar et al. [11] were able to prove the asymptotic behavior for the singular Yamabe problem in the flat case for any dimensions, for the general problem in the unit ball, F. Marques in [13] has proved the asymptotic behavior when the dimension is less or equal to five. For these dimensions he was able to overcome the lack of symmetry by constructing appropriate auxiliary functions. The problem for higher dimensions is still open.

Once that system (1) can be seen as a natural generalization of the singular Yamabe problem for an arbitrary metric, we need to put the same restrictions for the dimension. Furthermore, sometimes we will consider the following hypotheses on the potential

(H1) $-A$ is *cooperative*, that is, the components in the nondiagonal A_{ij} of A , $i \neq j$, are nonpositive;

(H2) In dimension $n = 5$, there exists a C^2 -function f such that

$$A(x) = f(x)Id_2 + O(|x|)$$

near the origin, where Id_2 is the identity matrix;

Under these assumptions we are able to prove our main theorem.

Theorem 0.3 *Assume $3 \leq n \leq 5$ and let \mathcal{U} be a positive solution for (1) in the punctured unit ball, and suppose that the potential A satisfies (H1) and (H2). If \mathcal{U} has a nonremovable singularity at 0 then there exists a Fowler-type solution \mathcal{U}_0 of (6) such that*

$$\mathcal{U}(x) = (1 + O(|x|^\alpha))\mathcal{U}_0(x)$$

as $x \rightarrow 0$, for some $\alpha > 0$.

The strategy of the proof of Theorem 0.3 relies on the properties of Fowler solutions. First, using the moving planes technique we prove the upper bound

$$|\mathcal{U}|(x) \leq c|x|^{\frac{2-n}{2}}. \tag{9}$$

As a consequence of this upper bound we can study *Pohozaev integrals* for solutions of (1), and use a Pohozaev-type identity to show that the invariant

$$P(\mathcal{U}) = \lim_{r \rightarrow \infty} P(r, \mathcal{U})$$

is well-defined, where

$$P(r, \mathcal{U}) = \int_{\partial B_r} \left(\frac{n-2}{2} \left\langle \mathcal{U}, \frac{\partial \mathcal{U}}{\partial \nu} \right\rangle - \frac{r}{2} |\nabla \mathcal{U}|^2 + r \left| \frac{\partial \mathcal{U}}{\partial \nu} \right|^2 + r \frac{(n-2)^2}{8} |\mathcal{U}|^{\frac{2n}{n-2}} \right) d\sigma.$$

Inspired by the works of L. Caffarelli, B. Gidas and J. Spruck in [3], we prove a removable singularity theorem, proving that this invariant is always nonpositive, and it is equal to zero if and only if the singularity is removable. As a consequence of this result we establish the lower bound

$$|\mathcal{U}|(x) \geq c|x|^{\frac{2-n}{2}}. \quad (10)$$

Using these bounds our goal is to use a scaling argument due to Leon Simon to prove Theorem 0.3. For this purpose, the study of the growth of what we call *Jacobi fields* for the limit system will be fundamental.

The thesis is organized as follows. In Chapter 1 we are going to do a overview on Fowler solutions and Jacobi fields. In Chapter 2 we will prove the radial symmetry for solutions to the limit system with intent to transform the system (6) in a ODE. Then we prove Theorem 0.2 and study the Jacobi fields associated to (4). In Chapter 3 we prove the upper and lower bounds for solutions of (4) and the asymptotic behavior for solutions of system (4). Finally in Chapter 4.3 we prove Theorem 0.3 assuming the hypothesis (H1) and (H2) on the potential A.

Notation and terminology

- $B_R(x)$ denotes the open ball of radius R and center x ;
- $B_R = B_R(0)$ denotes the open ball of radius R and center 0;
- χ_E denotes the characteristic function of a set $E \subset \mathbb{R}^n$, that is

$$\chi_E(x) = \begin{cases} 1 & \text{if } x \in E \\ 0 & \text{if } x \in \mathbb{R}^n \setminus E; \end{cases}$$

- $o(1)$ denotes a sequence which converges to zero;
- For $1 \leq p < \infty$, $L^p(B_1(0))^2 = L^p(B_1(0)) \times L^p(B_1(0))$ denotes the Lebesgue space with norm

$$\|(u, v)\|_p^p = \|u\|_p^p + \|v\|_p^p;$$

- $\mathbb{S}_+^{n-1} = \{x = (x_1, \dots, x_n) \in \mathbb{S}^{n-1} : x_i \geq 0 \quad \forall i\}$
- $\Delta_{\mathbb{S}^{n-1}}$ denotes the Laplacian in the unit sphere;
- If $\mathcal{U} = (u_1, u_2)$ then $|\mathcal{U}|^2 = \sum_{i=1}^2 u_i^2$, $|\nabla \mathcal{U}|^2 = \sum_{i=1}^2 |\nabla u_i|^2$ and $|\frac{\partial \mathcal{U}}{\partial \nu}|^2 = \sum_{i=1}^2 \frac{\partial u_i}{\partial \nu}^2$;
- g_{cyl}, g_{euc}

Chapter 1

Fowler solutions and Jacobi fields

As mentioned earlier, our main goal in this thesis is to show that the solutions of our system are asymptotic the solutions of the limit system. As these solutions will play a similar role to Fowler's solutions in the case of the singular Yamabe problem, we will briefly review the main properties of Fowler's solutions as well as what we call the Jacobi fields.

We say that a u is a *Fowler* or *Delaunay-type* solution if satisfies

$$\Delta u + \frac{n(n-2)}{4} u^{\frac{n+2}{n-2}} = 0 \quad \text{in } \mathbb{R}^n \setminus \{0\}, \quad (1.1)$$

and if it is singular around the origin. Using the Alexandrov reflection argument, the following result of Caffarelli, Gidas and Spruck in [3] characterizes these solutions.

Proposition 1.1 [3] *Let u be a positive solution of (1.1). If the origin is a non-removable singularity, then u is radially symmetric about the origin.*

Using that u is rotationally invariant the equation can be reduced to an ODE. Indeed, consider the diffeomorphism

$$\begin{aligned} \Phi : (\mathbb{R} \times \mathbb{S}^{n-1}, g_{cyl} = dt^2 + d\theta^2) &\rightarrow (\mathbb{R}^n \setminus \{0\}, g_{euc}) \\ \Phi(t, \theta) &= e^{-t}\theta. \end{aligned}$$

with inverse $\Phi^{-1}(x) = (-\log|x|, x|x|^{-1})$ and such that $\Phi^*g_{euc} = e^{-2t}g_{cyl}$. This diffeomorphism induces the classical change of variables from Fowler, given by

$$v(t, \theta) = |x|^{\frac{n-2}{2}} u(x). \quad (1.2)$$

where $t = -\log |x|$ and $\theta = x/|x|$.

Relative to this change of variables the equation (1.1) is equivalent to

$$v'' - \frac{(n-2)^2}{4}v + \frac{n(n-2)}{4}v^{\frac{n+2}{n-2}} = 0. \quad (1.3)$$

We analyse this equation by converting it into a system of first order equations

$$\begin{cases} v' = w \\ w' = \frac{(n-2)^2}{4}v - \frac{n(n-2)}{4}v^{\frac{n+2}{n-2}}. \end{cases}$$

whose Hamiltonian energy, given by

$$H(v, w) = w^2 - \frac{(n-2)^2}{4}v^2 + \frac{(n-2)^2}{4}v^{\frac{2n}{n-2}},$$

is constant along solutions of (1.3).

By examining the level curves of energy, we see that all positive solutions lie in the bounded set $\{H < 0\} \cap \{v > 0\}$.




Figure 1.1: Level sets

The basic properties of these solutions are summarized in the next proposition which is proved in [14].

Proposition 1.2 [14] *For any $H_0 \in (-(n-2)/n)^{n/2}(n-2)/2, 0)$, there exists a unique bounded solution of (1.3) satisfying $H(v, v') = H_0$, $v'(0) = 0$ and $v''(0) > 0$. This solution is periodic, and for all $t \in \mathbb{R}$ we have $v(t) \in (0, 1)$. This solution can be indexed by the parameter $\varepsilon = v(0) \in (0, ((n-2)/n)^{(n-2)/4})$, which is the smallest of the two values v assumes when $v'(0) = 0$. When $H_0 = -((n-2)/n)^{n/2}((n-2)/2)$ there is a unique bounded solution of (1.3), given by*

$$v(t) = \left(\frac{n-2}{n} \right)^{\frac{n-2}{4}}.$$

Finally, if v is a solution with $H_0 = 0$, then either $v(t) \equiv 0$ or $v(t) = (\cosh(t - t_0))^{\frac{2-n}{2}}$ for some $t_0 \in \mathbb{R}$.

1.1 The linearized equation

After introducing the notion of Fowler solutions, we describe some useful properties of the Jacobi fields associated to these solutions. The idea of using Jacobi fields in this context arose initially in the study of the asymptotic behavior of solutions to the conformally invariant semilinear equation

$$\Delta u + \frac{n(n-2)}{4} u^{\frac{n+2}{n-2}} = 0, \quad (1.4)$$

which using the cylindrical coordinates change is equivalent to

$$H(v) = \partial_t^2 v + \Delta_{\mathbb{S}^{n-1}} v - \frac{(n-2)^2}{4} v + \frac{n(n-2)}{4} v^{\frac{n+2}{n-2}} = 0. \quad (1.5)$$

Note that a solution of the equation (1.4) solves the singular Yamabe problem in the flat case.

Inspired by the work [15], in [11] the authors considered the linearization of the operator in (1.5) around a Fowler solution v_ε given by

$$L_\varepsilon(v) = \frac{\partial}{\partial t} \Big|_{t=0} H(v_\varepsilon + tv) = \partial_t^2 + \Delta_{\mathbb{S}^{n-1}} - \frac{(n-2)^2}{4} + \frac{n(n+2)}{4} v_\varepsilon^{\frac{4}{n-2}}. \quad (1.6)$$

Since the operator above has periodic coefficients, it can be studied using the classical Floquet theory or also by separation of variables, see [17]. Note that this operator can also be defined as the derivative at $s = 0$ of $H(U(s))$, where $U(s)$ is any one-parameter family of solutions with $U(0) = v_\varepsilon$, $U'(0) = v$.

This linearization was studied using results due to R. Mazzeo, D. Pollack and K. Uhlenbeck in [15] based on the spectral decomposition of the laplacian operator $\Delta_{\mathbb{S}^{n-1}}$. They were able to conclude that $\text{spec}(L_\varepsilon)$ is purely absolutely continuous, with no singular continuous or point spectrum.

If $\{\lambda_k, \mathcal{X}_k(\theta)\}$ is the eingendata of $\Delta_{\mathbb{S}^{n-1}}$, using the convention that these eigenvalues are listed with multiplicity, we can write $\lambda_0 = 0$, $\lambda_1 = \dots = \lambda_n = n-1$, $\lambda_{n+1} = 2n$ and so on. Hence the linearized operator could be decomposed into many ordinary differential operators given by

$$L_{\varepsilon,k} = \partial_t^2 + \left(\frac{n(n+2)}{4} v_\varepsilon^{\frac{4}{n-2}} - \frac{(n-2)^2}{4} - \lambda_k \right). \quad (1.7)$$

What we call **Jacobi fields** are the elements in the kernel of the linearized operator, that is, the functions ψ which satisfy $L_\varepsilon \psi = 0$. The properties of these fields are important for the spectral analysis of the linearized near origin.

It is worth noting that considering the linearization of the equation around a Fowler solution is a method used not only to study the asymptotic behavior of the solutions. In fact, the analysis of the linearized operator together with fixed point theorems are the main tools used in the proof of the existence of the singular Yamabe problem, under certain hypotheses, as applied in the works of [20] and [14].

In order to study the properties of the Jacobi fields using the decomposition of the laplacian in terms of the eigenvalues, it is sufficient to consider solutions to the induced problems $L_{\varepsilon,k}(\psi_k) = 0$.

It is common, in geometric problems, that some solutions to low values of k , not necessarily all, are obtained as derivatives of one-parameter families of solutions to (1.5). For the first eigenvalue $\lambda_0 = 0$, if we consider the families of solutions to (1.5) given by

$$T \rightarrow v_\varepsilon(t + T) \quad \text{and} \quad \varepsilon \rightarrow v_\varepsilon(t), \quad (1.8)$$

and then differentiate these families with respect to the parameters to obtain solutions to (1.7) corresponding to λ_0 and given by

$$\psi_{\varepsilon,0}^+(t) = \frac{d}{dT}v_\varepsilon(t + T) = v'_\varepsilon(t + T), \quad \psi_{\varepsilon,0}^-(t) = \frac{d}{d\varepsilon}v_\varepsilon(t),$$

which are linearly independent Jacobi fields with linearly and periodic growth, respectively. They also proved that these two functions form a basis, for each ε , for all temperate solutions of $L_\varepsilon\psi = 0$ on the cylinder. Any solution of this equation must grow exponentially in one direction or the other.

Using a similar construction for $\lambda_k = n - 1$, they were able to build linearly independent solutions $\psi_{\varepsilon,k}^\pm$ that are exponentially increasing and decreasing. Finally when $k > n + 1$, we know that the term of zero order of the above operator is negative, since $\lambda_k \geq 2n$ and $v_\varepsilon < 1$ which imply that $L_{\varepsilon,k}$ satisfies the maximum principle.

Chapter 2

Classification result for the Limit system

The main goal in this chapter is to study some properties of singular nonnegative solutions $\mathcal{U} = (u_1, u_2)$ to the following system

$$\Delta u_i + \frac{n(n-2)}{4} |\mathcal{U}|^{\frac{4}{n-2}} u_i = 0 \quad \text{in } \mathbb{R}^n \setminus \{0\},$$

where $|\mathcal{U}|^2 = \sum_{i=1}^2 u_i^2$ and Δ denotes the euclidian laplacian. Similarly to the definition for the system (1), we say that \mathcal{U} is a nonnegative solution if each coordinate u_i is nonnegative and we say that it is singular if the following holds

$$\lim_{|x| \rightarrow 0} |\mathcal{U}|(x) = +\infty. \tag{2.1}$$

It may happen that only one of the coordinates blows up at the singularity.

The above system will be important since its solutions play a similar role to Fowler solutions in the singular Yamabe problem.

One of our main result in this chapter will be the characterization of the nonnegative singular solutions to the limit system. We will show that all such solutions are Fowler solution times a vector in the unit sphere with nonnegative coordinates (see Theorem 2.10 below). Another important result will be describe when the singularity can be removed in terms of an *Pohozaev invariant* inspired by the works of Caffarelli, Gidas and Spruck [3] and R. Fowler [9]. We will present two distinct ways of proving this result.

2.1 Radial Symmetry

In the first step in this direction we will be using a “measure theoretic” variation of the Alexandrov reflection technique as developed by Gidas, Ni and Nirenberg [10], to prove the radial symmetry of nonnegative singular solutions. To proving this result, we need of the following result from Chen and Lin [4], which ensures that the maximum principle holds for superharmonic functions with isolated singularities.

Lemma 2.1 [4] *Let E be a smooth bounded domain in \mathbb{R}^n , and Z a compact set in \mathbb{R}^n with $\text{Cap}(Z) = 0$. Assume that $v(x)$, $h(x)$ are nonnegative continuous functions in $\bar{E} \setminus Z$ and satisfy*

$$\Delta v(x) + h(x) \leq 0 \quad \text{in} \quad E \setminus Z,$$

in the distributional sense. Then

$$v(x) \geq \int_E G(x, y) h(y) dy + \int_{\partial E} \frac{\partial G(x, y)}{\partial \nu} v(y) ds_y$$

for $x \in E \setminus Z$, where $G(x, y)$ is the Green function of $-\Delta$ in E with the Dirichlet boundary condition. In particular, we have $v(x) \geq \inf_{\partial E \setminus Z} v$.

Theorem 2.2 (Radial Symmetry) *Let $n \geq 3$ and $\mathcal{U} = (u_1, u_2)$ be a nonnegative C^2 singular solution of the nonlinear elliptic system (6). Then \mathcal{U} is radially symmetric about the origin.*

Proof. Without loss of generality, we may assume that $\lim_{|x| \rightarrow 0} u_1(x) = +\infty$. Now, we will use the moving planes method.

Fix an arbitrary $z \neq 0$ in \mathbb{R}^n . Define the Kelvin transform

$$U_i(x) = \frac{1}{|x|^{n-2}} u_i \left(z + \frac{x}{|x|^2} \right),$$

for $i \in \{1, 2\}$. By Caffarelli et al. [3], the Kelvin transform U_i has the asymptotic expansion at infinity [3, Lemma 2.3]. Since 0 is a nonremovable singularity for u_1 we can see that U_1 is singular at 0 and $z_0 = -z/|z|^2$, and U_2 is singular at 0. Moreover, the Kelvin transform satisfies

$$-\Delta U_i = \frac{n(n-2)}{4} |\mathbf{U}|^{\frac{4}{n-2}} U_i \quad \text{in} \quad \mathbb{R}^n \setminus \{0, z_0\}, \quad (2.2)$$

where $|\mathbf{U}|^2 = \sum_{i=1}^2 U_i^2$. It is easy to see that

$$U_i(x) = O(|x|^{2-n}) \quad \text{as} \quad |x| \rightarrow \infty$$

and that each U_i has the following harmonic asymptotic expansion at infinity

$$U_i(x) = a_{i0}|x|^{2-n} + a_{ij}x_j|x|^{-n} + O(|x|^{-n}),$$

$$\partial_{x_j}U_i = (2-n)a_{i0}x_j|x|^{-n} + O(|x|^{-n})$$

and

$$\partial_{x_k}\partial_{x_j}U_i = O(|x|^{-n}),$$

where $a_{i0} = u_i(z)$ and $a_{ij} = \partial_{y_j}u_i(z)$.

We will show that U_i are axisymmetric with respect to the axis going through 0 and z . Choose a reflection direction orthogonal to this axis and assume without loss of generality that it is equal to the positive x_n direction $(0, \dots, 0, 1)$. For $\lambda > 0$ let $\Sigma_\lambda := \{x \in \mathbb{R}^n : x_n > \lambda\}$ and consider the reflection

$$x = (x_1, \dots, x_{n-1}, x_n) \in \Sigma_\lambda \mapsto x_\lambda = (x_1, \dots, x_{n-1}, 2\lambda - x_n).$$

Using again the asymptotic expansion at infinity for the Kelvin transform from [3], there exist positive constants $\bar{\lambda} > 10$ and $R > |z_0| + 10$ such that for any $\lambda \geq \bar{\lambda}$ we have, for each coordinate

$$U_i(x) < U_i(x_\lambda), \text{ for } x \in \Sigma_\lambda \text{ and } |x_\lambda| > R.$$

By Lemma 2.1, there exists a constant $C > 0$ such that

$$U_i(x) \geq C, \quad \text{for } x \in \bar{B}_R \setminus \{0, z_0\}.$$

Since $U_i(x) \rightarrow 0$ as $|x| \rightarrow +\infty$, there exists $\lambda_0 > \bar{\lambda}$ such that when $|x| \geq 2\lambda_0 - R$ it holds $U_i(x) < C$. On the other hand if $x \in \Sigma_\lambda$ and $x_\lambda \in \bar{B}_R$ then $x \notin B_{2\lambda-R}$. Thus for every $\lambda \geq \lambda_0$ we have

$$U_i(x) \leq U_i(x_\lambda), \text{ for all } x \in \Sigma_\lambda \text{ s.t. } x_\lambda \notin \{0, z_0\}. \quad (2.3)$$

Let

$$\lambda^* := \inf\{\tilde{\lambda} > 0 \mid (2.3) \text{ holds for all } \lambda \geq \tilde{\lambda}\}.$$

It suffices to show that $\lambda^* = 0$. Indeed, this proves that U_i is axisymmetric with respect to the axis going through 0 and z and since z is arbitrary, each u_i is radially symmetric about the origin.

Suppose by contradiction that $\lambda^* > 0$. Then (2.3) holds for $\lambda = \lambda^*$. Since U_1 blows-up at z_0 we see that U_1 cannot be invariant by the reflection x_{λ^*} . Then by applying the maximum principle to $U_1(x_{\lambda^*}) - U_1(x)$ and $U_2(x_{\lambda^*}) - U_2(x)$ in (2.2), we conclude that

$$U_i(x) < U_i(x_{\lambda}), \text{ for } x \in \Sigma_{\lambda} \text{ and } x_{\lambda} \notin \{0, z_0\}. \quad (2.4)$$

Then applying the Hopf boundary lemma for $x \in \partial\Sigma_{\lambda^*}$ we obtain

$$\partial_{x_n}(U_i(x_{\lambda^*}) - U_i(x)) = -2\partial_{x_n}U_i(x) > 0. \quad (2.5)$$

Now choose sequences $\lambda_j \nearrow \lambda^*$ and $x^j \in \Sigma_{\lambda_j}$ such that $U_1(x_{\lambda_j}^j) < U_1(x^j)$. By [3, Lemma 2.4] (the plane $x_n = 0$ there corresponds to $x_n = \lambda^*$ here), we conclude that the sequence $|x^j|$ is bounded. Hence passing to a subsequence we may assume that $x_j \rightarrow \bar{x} \in \Sigma_{\lambda^*}^-$ with $U_1(\bar{x}_{\lambda^*}) \leq U_1(\bar{x})$. By (2.4) we know that $\bar{x} \in \partial\Sigma_{\lambda^*}$ and then $\partial_{x_n}U_1(\bar{x}) \geq 0$, a contradiction with (2.5). Therefore, $\lambda^* = 0$. ■

Once we proved the radial symmetry of the solutions we may use the classical change of variables from Fowler to approach our problems using ODE analysis. This is the subject of the next section.

2.2 ODE Analysis

If $\mathcal{U} = (u_1, u_2)$ is a solution of the system, from (2.2) we may assume that each $u_i(x) = u_i(|x|)$. So using the classical change of variables from Fowler (1.2), it follows that $V = (v_1, v_2)$ satisfies the ordinary differential equation

$$\begin{cases} v_i'' - \delta^2 v_i + \frac{n(n-2)}{4}|V|^{\frac{4}{n-2}}v_i = 0, & t \in \mathbb{R} \\ v_i \geq 0, & v_i \in C^2(\mathbb{R}). \end{cases} \quad (2.6)$$

where $\delta = (n-2)/2$. Define

$$\Psi(t) := \frac{1}{2}(|V'|^2 - \delta^2|V|^2)(t) + \frac{\delta^2}{2}|V|^{\frac{2n}{n-2}}(t).$$

Then $\Psi'(t) \equiv 0$, namely $\Psi(t)$ is a constant K . As a consequence it is easy to see that $v_i, |v_i'|$ are bounded. Indeed, if we suppose that there exists a sequence $t_n \rightarrow +\infty$ such that $|V(t_n)| \rightarrow +\infty$, we have

$$\frac{1}{2} \left[\left(\frac{|V'(t_n)|}{|V(t_n)|} \right)^2 - \delta^2 \right] + \frac{\delta^2}{2}|V(t_n)|^{\frac{4}{n-2}} = \frac{K}{|V(t_n)|^2}$$

which is a contradiction. Similarly, we conclude that $|v'_i|$ is bounded.

The first consequence of the radial symmetry of solutions is the fact that each coordinate u_i satisfies the following dichotomy: either it is strictly positive, or it vanishes identically.

Lemma 2.3 *Let \mathcal{U} a nonnegative singular solution of the limit system (6). If there exist $y \in \mathbb{R}^n \setminus \{0\}$ such that $u_i(y) = 0$, for some $i \in \{1, 2\}$ then $u_i \equiv 0$.*

Proof. We already know that our coordinates are radially symmetric. Suppose that there exist $y, z \in \mathbb{R}^n \setminus \{0\}$, with $|z| > |y| = r$, $u_i(y) = u_i(r) = 0$ and $u_i(|z|) > 0$. By the maximum principle of Chen and Lin in [4], we know that

$$u_i(x) \geq \inf_{\partial B_{|z|}} u_i = u_i(|z|) > 0, \quad \forall x \in B_{|z|}(0) \setminus \{0\}$$

which contradicts the fact that $u_i(y) = 0$. Hence, if there exists a point y such that $u_i(y) = 0$ then u_i must be zero for all z with $|z| \geq |y|$.

However, if we consider the change of variables of Fowler, we know that v_i is a solution for the ODE system (2.6), and by the uniqueness of solutions to ODE systems, it would be identically zero. ■

In order to prove the classification result, we will need the following technical results

Lemma 2.4 *Fix $i \in \{1, 2\}$. If v_i has a limit C_i at $+\infty$ or $-\infty$, then $C_i \leq \left(\frac{n-2}{n}\right)^{\frac{n-2}{4}}$.*

Proof. Assume $\lim_{t \rightarrow +\infty} v_1(t) = C_1$ where $C_1 > 0$. For each coordinate consider a sequence of translates $v_{i,k}(t) = v_i(t + k)$. By (2.6), up to passing to a subsequence, we may assume that $v_{1,k} \rightarrow C_1$ and $v_{2,k}^2 \rightarrow w \geq 0$ uniformly in compacts with respect to the C^2 topology. Thus $-\delta^2 C_1 + \frac{n(n-2)}{4}(C_1^2 + w)^{\frac{2}{n-2}} C_1 = 0$ which implies that $C_1 \leq \left(\frac{n-2}{n}\right)^{\frac{n-2}{4}}$. ■

For each i such that the coordinate u_i is strictly positive, we define the auxiliary functions $f_i : \mathbb{R} \rightarrow \mathbb{R}$ by

$$f_i(t) := -\frac{1}{2}|v'_i(t)|^2 + \frac{\delta^2}{2}v_i(t)^2 - \frac{\delta^2}{2}v_i(t)^{\frac{2n}{n-2}}. \quad (2.7)$$

By (2.6), we have $f'_i(t) = \frac{\delta^2}{2}(|V|^{\frac{4}{n-2}}(t) - v_i(t)^{\frac{4}{n-2}})v_i(t)v'_i(t)$. That is, the functions f_i and v_i have exactly the same monotonicity. This property is very important and will be used in the sequel.

Lemma 2.5

$$v_i(t) < 1, \quad \forall t \in \mathbb{R} \text{ and } i = 1, 2.$$

Proof. Suppose by contradiction that the result is not true. By the Lemma 2.4, there exists a local maximum point $t_0 \in \mathbb{R}$ of v_1 such that $v_1(t_0) \geq 1$. This implies that

$$f_1(t_0) = \frac{\delta^2}{2} v_1^2(t_0) - \frac{\delta^2}{2} v_1(t_0)^{\frac{2n}{n-2}} \leq 0.$$

We claim that there exists $t_1 > t_0$ such that $v_1'(t) < 0$ for $t \in (t_0, t_1)$ and $v_1'(t_1) = 0$. Otherwise it would follow from Lemma 2.4 that $v_1 \downarrow C_1 \leq ((n-2)/n)^{\frac{n-2}{4}}$ and $f_1(t) \downarrow -C < 0$ as $t \uparrow +\infty$. Then we have that $v_1'(t) \rightarrow 0$ as $t \uparrow +\infty$ and so

$$0 > -C = \frac{\delta^2}{2} C_1^2 - \frac{\delta^2}{2} C_1^{\frac{2n}{n-2}} > 0,$$

a contradiction. Hence there exists such t_1 and it satisfies $v_1''(t_1) \geq 0$. On the other hand

$$f_1(t_1) = \frac{\delta^2}{2} v_1(t_1)^2 - \frac{\delta^2}{2} v_1(t_1)^{\frac{2n}{n-2}} < f_1(t_0) \leq 0,$$

namely $v_1(t_1) > 1$ which implies from (2.6) that $v_1''(t_1) < 0$, a contradiction. ■

Lemma 2.6 $f_i(t) > 0$ and so $|v_i'(t)| < \delta v_i(t)$ for all $t \in \mathbb{R}$ and $i = 1, 2$.

Proof. Suppose $f_1(s) \leq 0$ for some $s \in \mathbb{R}$. If $f_1'(s) = 0$, then $v_1'(s) = 0$, which implies by (2.7) that $v_1(s) \geq 1$, a contradiction with Lemma 2.5. So we may assume, without loss of generality, that $f_1'(s) > 0$, namely $v_1'(s) > 0$. If $v_1'(t) > 0$ for all $t \leq s$, then $v_1(t) \downarrow C_1 \leq \left(\frac{n-2}{n}\right)^{\frac{4}{n-2}}$ and $f_1(t) \downarrow -C < 0$ as $t \downarrow -\infty$, and we get a contradiction as in the proof of Lemma 2.5. So there exists $t_0 < s$ such that $v_1'(t) > 0$ for $t \in (t_0, s]$ and $v_1'(t_0) = 0$. Similarly we get $v_1(t_0) > 1$, a contradiction with Lemma 2.5. ■

2.3 Removable singularity result

We now turn to a discussion of the existence and specific form of a family of homological integral invariants of solutions of system (1). These invariants were discovered in their simplest form by Pohozaev and was later put into a general Riemannian setting in the work of R. Schoen [19]. As we are working in a ball in the euclidian space, we can use a simple definition.

For $r > 0$, define the Pohozaev integral

$$P(r, \mathcal{U}) = \int_{\partial B_r} \left(\frac{n-2}{2} \left\langle \mathcal{U}, \frac{\partial \mathcal{U}}{\partial \nu} \right\rangle - \frac{r}{2} |\nabla \mathcal{U}|^2 + r \left| \frac{\partial \mathcal{U}}{\partial \nu} \right|^2 + r \frac{(n-2)^2}{8} |\mathcal{U}|^{\frac{2n}{n-2}} \right) d\sigma \quad (2.8)$$

where ν is the unit outer normal of ∂B_r . Multiplying each equation of (6) by $x \cdot \nabla u_i$ and integrating over $B_r \setminus B_s$ we verify for all $r, s > 0$ that

Lemma 2.7 (Pohozaev Identity) *For $0 < s \leq r$, we have*

$$P(r, \mathcal{U}) = P(s, \mathcal{U}).$$

Proof. Adding and subtracting Δu_i in each system equation, we have

$$\Delta u_i + \frac{n(n-2)}{4} |\mathcal{U}|^{\frac{4}{n-2}} u_i = 0 \quad (2.9)$$

in the punctured ball $\Omega = B_1^n(0) \setminus \{0\}$. Multiplying both sides by $x \cdot \nabla u_i$ and integrating over $B_r \setminus B_s$ we have

$$-\int_{B_r \setminus B_s} x \cdot \nabla u_i \left(\Delta u_i + \frac{n(n-2)}{4} |\mathcal{U}|^{\frac{4}{n-2}} u_i \right) = 0$$

First using integration by parts we get

$$\begin{aligned} \int_{B_r \setminus B_s} x \cdot \nabla u_i \Delta u_i &= \frac{n-2}{2} \int_{B_r \setminus B_s} |\nabla u_i|^2 - \frac{r}{2} \int_{\partial B_r} |\nabla u_i|^2 + \frac{s}{2} \int_{\partial B_s} |\nabla u_i|^2 \\ &+ r \int_{\partial B_r} \left| \frac{\partial u_i}{\partial \nu} \right|^2 - s \int_{\partial B_s} \left| \frac{\partial u_i}{\partial \nu} \right|^2. \end{aligned} \quad (2.10)$$

On the other hand, multiplying each equation for u_i and integrating over $B_r \setminus B_s$ we have

$$\begin{aligned} 0 &= - \int_{B_r \setminus B_s} u_i \left(\Delta u_i + \frac{n(n-2)}{4} |\mathcal{U}|^{\frac{4}{n-2}} u_i \right) = \\ &\int_{B_r \setminus B_s} |\nabla u_i|^2 - \int_{\partial B_r} u_i \frac{\partial u_i}{\partial \nu} + \int_{\partial B_s} u_i \frac{\partial u_i}{\partial \nu} - \frac{n(n-2)}{4} \int_{B_r \setminus B_s} |\mathcal{U}|^{\frac{4}{n-2}} u_i^2 \end{aligned}$$

which implies that

$$\int_{B_r \setminus B_s} |\nabla u_i|^2 = \int_{\partial B_r} u_i \frac{\partial u_i}{\partial \nu} - \int_{\partial B_s} u_i \frac{\partial u_i}{\partial \nu} + \frac{n(n-2)}{4} \int_{B_r \setminus B_s} |\mathcal{U}|^{\frac{4}{n-2}} u_i^2 \quad (2.11)$$

Replacing (4.27) in (2.10), we conclude that

$$\begin{aligned} &\int_{\partial B_r} \left[\frac{n-2}{2} u_i \frac{\partial u_i}{\partial \nu} - \frac{r}{2} |\nabla u_i|^2 + r \left| \frac{\partial u_i}{\partial \nu} \right|^2 \right] - \int_{\partial B_s} \left[\frac{n-2}{2} u_i \frac{\partial u_i}{\partial \nu} - \frac{s}{2} |\nabla u_i|^2 + s \left| \frac{\partial u_i}{\partial \nu} \right|^2 \right] \\ &= - \frac{n(n-2)^2}{8} \int_{B_r \setminus B_s} |\mathcal{U}|^{\frac{2n}{n-2}} - \frac{n(n-2)}{4} \int_{B_r \setminus B_s} (x \cdot \nabla u_i |\mathcal{U}|^{\frac{4}{n-2}} u_i). \end{aligned}$$

Note also that, using $\frac{n-2}{2n}\partial_k(|\mathcal{U}|^{\frac{2n}{n-2}}) = \sum_i |\mathcal{U}|^{\frac{4}{n-2}} u_i \partial_k u_i$, we obtain

$$\begin{aligned} - \sum_i \int_{B_r \setminus B_s} x \cdot \nabla u_i |\mathcal{U}|^{\frac{4}{n-2}} u_i &= \frac{n-2}{2} \int_{B_r \setminus B_s} |\mathcal{U}|^{\frac{2n}{n-2}} - r \frac{n-2}{2n} \int_{\partial B_r} |\mathcal{U}|^{\frac{2n}{n-2}} \\ &\quad + s \frac{n-2}{2n} \int_{\partial B_s} |\mathcal{U}|^{\frac{2n}{n-2}}, \end{aligned}$$

which finishes the proof.

■

From the above result we know that $P(r, \mathcal{U})$ is a constant independent of r that we denote by $P(\mathcal{U})$. We will prove that

Theorem 2.8 *Let \mathcal{U} be a positive solution of (6). Then $P(\mathcal{U}) \leq 0$. Moreover, $P(\mathcal{U}) = 0$ if and only if each coordinate u_i is smooth at 0.*

Proof. Our strategy to prove the result about removable singularity will be suppose that $P(\mathcal{U}) \geq 0$, to prove that necessarily we have $P(\mathcal{U}) = 0$ and then that all the components of the solution are smooth.

Suppose that $P(\mathcal{U}) \geq 0$. We want to prove that $P(\mathcal{U}) = 0$ and $u_i \in C^2(\mathbb{R}^n)$. Recall that $K \geq 0$. Then we have,

$$0 < \sum_{i=1}^2 f_i(t) = \frac{\delta^2}{2} |V|^{\frac{2n}{n-2}} - \sum_{i=1}^2 \frac{\delta^2}{2} v_i(t)^{\frac{2n}{n-2}} - K \leq \frac{\delta^2}{2} |V|^{\frac{2n}{n-2}} - \sum_{i=1}^2 \frac{\delta^2}{2} v_i(t)^{\frac{2n}{n-2}} \quad (2.12)$$

According to the Lemma 2.3, we can consider two different situations. If one of the coordinates for the solution vanishes, we have a contradiction from the above expression. Otherwise, Lemma 2.4 implies the existence of $s \in \mathbb{R}$ such that $v'_1(s) = 0$. Let $s \in \mathbb{R}$ be any a point such that $v'_1(s) = 0$. Then, by Lemma 2.6 and (2.12),

$$f_1(s) = \frac{\delta^2}{2} v_1(s)^2 - \frac{\delta^2}{2} v_i(s)^{\frac{2n}{n-2}} < \frac{\delta^2}{2} |V|^{\frac{2n}{n-2}} - \sum_{i=1}^2 \frac{\delta^2}{2} v_i(s)^{\frac{2n}{n-2}}.$$

Note that in the right side of the equation, we will only have terms that are multiples of $v_1(s)^2$. So dividing everything by $v_1(s)^2$ and estimating the terms that remain, we obtain by a direct computation

$$\frac{|V|^{\frac{2n}{n-2}} - \sum_{i=1}^2 v_i(s)^{\frac{2n}{n-2}}}{v_i(s)^2} + v_i(s)^{\frac{4}{n-2}} \leq C |V|^{\frac{4}{n-2}},$$

where $C = \frac{2^*}{2}$. Then

$$\delta^2 < \frac{n(n-2)}{4} |V|^{\frac{4}{n-2}}. \quad (2.13)$$

This inequality implies that

$$v_1''(s) = \left(\delta^2 - \frac{n(n-2)}{4} |V|^{\frac{4}{n-2}} \right) v_1(s) < 0.$$

That is, s must be a local maximum point of v_1 , and v_1 has no local minimum points. Therefore, v_1 has a unique maximum point t_1 , $v_1'(t) > 0$ for $t < t_1$ and $v_1'(t) < 0$ for $t > t_1$. Similarly, v_2 has a unique maximum point t_2 , $v_2'(t) > 0$ for $t < t_2$. Lemma 2.4 gives that $v_2 \downarrow 0, \forall i$ as $t \uparrow +\infty$. This, together with Lemma 2.6, yields $K = 0$, namely $P(\mathcal{U}) = 0$. Let $v = \sum_{i=1}^2 v_i$ and $y(t) = v'(t) + \delta v(t)$ then $v'(t) < 0$ for all $t > \max_{1 \leq i \leq 2} t_i$ and

$$y' - \delta y = v'' - \delta^2 v = -\frac{n(n-2)}{4} |V|^{\frac{4}{n-2}} v \geq -C v^{\frac{n+2}{n-2}}.$$

Since $y > 0$ is bounded, it is easy to prove that

$$v'(t) + \delta v(t) = y(t) \leq \frac{C}{\delta} v^{\frac{n+2}{n-2}}, \quad \forall t > \max_{1 \leq i \leq 2} t_i$$

and then $(e^{\delta t} v(t))^{2-2^*} - \frac{C}{\delta} e^{(2-2^*)\delta t}$ is strictly increasing for $t > \max_{1 \leq i \leq 2} t_i$. Combining this with $v(t) \downarrow 0$ as $t \uparrow +\infty$ we get $e^{\delta t} v(t) \leq C$ uniformly for $t > 0$ large. That is $\sum_{i=1}^2 u_i(r) \leq C$ uniformly for $r > 0$ small. Therefore, for standard elliptic regularity theory, $u_i \in C^2(\mathbb{R}^n), \forall i$, which ends the proof.

■

2.4 Classification Result

In the beggining of this chapter, we mentioned that the solutions of our limit system play a similar role to Fowler's solutions in the singular Yamabe problem.

On the other hand, O. Druet, E. Hebey and J. Vétois in [6], [7] proved the following characterization result for entire solutions in \mathbb{R}^n for the limit system

Proposition 2.9 ([7]) *Let \mathcal{U} be a nonnegative C^2 -solution of*

$$\Delta u_i + |\mathcal{U}|^{\frac{4}{n-2}} u_i = 0 \quad \text{in } \mathbb{R}^n.$$

Then there exist $a \in \mathbb{R}^n$, $\mu > 0$ and $\Lambda \in \mathbb{S}_+^1$, such that

$$\mathcal{U}(x) = \left(\frac{\mu}{\mu^2 + \frac{|x-a|^2}{n(n-2)}} \right)^{\frac{n-2}{2}} \Lambda,$$

for all $x \in \mathbb{R}^n$, where \mathbb{S}_+^1 consists of the elements (Λ_1, Λ_2) in the unit sphere such that each coordinate is nonnegative.

Inspired by their result, we may ask wheter a similar description holds in the singular case. Indeed, we will prove that every nonnegative singular solution of (6) it is a Fowler solution times a vector in the unit sphere with nonnegative coordinates.

Theorem 2.10 *Suppose that \mathcal{U}_0 is a nonnegative singular solution for (6). Then there exists a $\Lambda \in \mathbb{S}_+^1$ and u_0 a Fowler solution such that*

$$\mathcal{U}_0 = u_0 \Lambda.$$

where $\mathbb{S}_+^1 = \{x \in \mathbb{S}^1 : x_i \geq 0\}$.

Proof. We will prove the result in the case where v_1 and v_2 are positive solutions since otherwise the result follows directly from Lemma 2.3.

Let $\mathcal{V} = (v_1, v_2)$ be the solution for (2.6) obtained from \mathcal{U}_0 after the change of variables of Fowler. We know that it satisfies

$$v_i'' - \delta^2 v_i + \frac{n(n-2)}{4} |\mathcal{V}|^{\frac{4}{n-2}} v_i = 0, \quad t \in \mathbb{R}.$$

From this, we can note that

$$v_1''(t)v_2(t) = v_1(t)v_2''(t),$$

which implies

$$v_1'(t)v_2(t) - v_1(t)v_2'(t) = c,$$

for all $t \in \mathbb{R}$, where c is a constant.

Suppose that $c \neq 0$. Whitout loss of generality we can assume that $c > 0$. Then by Lemma 2.4 we obtain

$$\left(\frac{v_1}{v_2}\right)'(t) = \frac{c}{v_2^2(t)} \geq c \quad (2.14)$$

which is a contradiction, since by (2.14) the quocient would assume negative values.

Therefore we conclude that

$$\left(\frac{v_1}{v_2}\right)'(t) = 0$$

and so $v_1/v_2 = \eta$ is constant. It follows that

$$\mathcal{V} = (v_1, v_2) = (\eta v_2, v_2) = (\eta, 1)v_2 = \sqrt{1 + \eta^2} v_2 \left(\frac{\eta}{\sqrt{1 + \eta^2}}, \frac{1}{\sqrt{1 + \eta^2}} \right)$$

and a direct computation shows that $v_0 = \sqrt{1 + \eta^2} v_2$ is a Fowler solution.

■

Although the proof above is quite simple, the result will be extremely important in the study of the Jacobi fields since will simplify the linearized system, and it also allows us to use all of the properties from Fowler solutions that we described previously. Inspired by this result, we will call the solutions of the system (6) of *Fowler-type solutions* or *Delaunay-type solutions*.

Let us mention some important consequences of the theorem. We observed in the introduction of this section that the definition of singular solution thus not imply that both solutions blows up at the origin but, as consequence of Theorem (2.10), we can see that this is exactly what happens when both coordinates are strictly positive.

Corollary 2.11 *Suppose that $\mathcal{U} = (u_1, u_2)$ is a solution for (6), such that both coordinates are strictly positive. Then $\lim_{|x| \rightarrow 0} u_i(x) = +\infty$, for $i = 1, 2$.*

Our second direct consequence will be the following asymptotic behavior

Corollary 2.12 *If \mathcal{U} is a nonnegative singular solution to the system (6), then \mathcal{U} is an admissible solution, that is, there exists positive constants c_1, c_2 such that*

$$c_1|x|^{\frac{2-n}{2}} \leq |\mathcal{U}|(x) \leq c_2|x|^{\frac{2-n}{2}}. \quad (2.15)$$

Besides note that by the classification result, there exists a Fowler solution u_0 and an unit vector $\Lambda = (\Lambda_1, \Lambda_2)$ with nonnegative coordinates, such that $\mathcal{U} = u_0\Lambda$. Consequently $P(\mathcal{U})$ coincides with the Pohozaev invariant of u_0 as defined in [3] which implies that the removable singularity result (2.8) proved in one of the previous sections follow as a direct consequence of Theorem (2.10).

2.5 Jacobi Fields for the Limit System

Carrying out the study intended in the following chapters, in this last section we will study what we call the Jacobi fields for the limit system.

In the study of the asymptotic behavior of the singular Yamabe problem in the punctured ball for flat and non flat case, N. Korevaar et al. in [11] and F. Marques [13] respectively, used the previously known growth properties of Jacobi fields as a tool to obtain information about the singular problem.

In order to do a similar analysis we are going to linearize solutions of system around the Fowler-type solutions that we characterized in the previous subsection. Consider the operator in cylindrical coordinates $H(\mathcal{V}) = (H^1(\mathcal{V}), H^2(\mathcal{V}))$ where

$$H^i(\mathcal{V}) = \partial_t^2 v_i + \Delta_{\mathbb{S}^{n-1}} v_i - \frac{(n-2)^2}{4} v_i + \frac{n(n-2)}{4} |\mathcal{V}|^{\frac{4}{n-2}} v_i. \quad (2.16)$$

When we linearize this operator on a solution $\mathcal{V}_0 = (v_{0,1}, v_{0,2})$, we obtain that the linearized operator is given by $\mathcal{L}(\phi) = (\mathcal{L}^1(\phi), \mathcal{L}^2(\phi))$ where

$$\begin{aligned} \mathcal{L}^i(\phi) &= \left. \frac{\partial}{\partial t} \right|_{t=0} H^i(\mathcal{V}_0 + t\phi) \\ &= \partial_t^2 \phi_i + \Delta_{\mathbb{S}^{n-1}} \phi_i - \frac{(n-2)^2}{4} \phi_i + \frac{n(n-2)}{4} \left(\frac{4}{n-2} |\mathcal{V}_0|^{2^*-4} \langle \mathcal{V}_0, \phi \rangle v_{0,i} + |\mathcal{V}_0|^{\frac{4}{n-2}} \phi_i \right). \end{aligned}$$

However from the characterization result obtained in Theorem 2.10, we know that there exist a Fowler solution v_ε and a vector in the unit sphere $\Lambda = (\Lambda_1, \Lambda_2)$ with positive coordinates such that $V_0 = v_\varepsilon \Lambda$. Using this fact, our linearized system can be simplified and written as follows

$$\mathcal{L}_\varepsilon^i(\phi) = \partial_t^2 \phi_i + \Delta_{\mathbb{S}^{n-1}} \phi_i - \frac{(n-2)^2}{4} \phi_i + n\Lambda_i \langle \Lambda, \phi \rangle v_\varepsilon^{\frac{4}{n-2}} + \frac{n(n+2)}{4} v_\varepsilon^{\frac{4}{n-2}} \phi_i. \quad (2.17)$$

We will focus now in the study of the **Jacobi fields** for the system, that is, solutions of the linear ODE system $L_\varepsilon(\mathcal{V}) = 0$.

Similarly to the case of a single equation, we can consider the spectral decomposition of the operator $\Delta_{\mathbb{S}^{n-1}}$ writing $\{\lambda_k, \mathcal{X}_k(\theta)\}$ for its eigendata and if $\phi = (\phi_1, \phi_2)$ is a solution of the system $\mathcal{L}_\varepsilon^i(\phi) = 0$, then for each coordinate

$$\phi_i(t) = \sum v_{i,k}(t) \mathcal{X}_k(\theta)$$

where $v_{i,k}$ satisfies

$$\mathcal{L}_{\varepsilon,k} \phi_{i,k} + n\Lambda_i \langle \Lambda, \phi_k \rangle v_\varepsilon^{\frac{4}{n-2}} + \frac{n(n-2)}{4} v_\varepsilon^{\frac{4}{n-2}} \phi_i = 0$$

and $\mathcal{L}_{\varepsilon,k} = \partial_t^2 - (\lambda_k + \frac{(n-2)^2}{4})$.

We are going to obtain the Jacobi fields for the first eigenvalue and we will see that the construction for higher eigenvalues is similar.

In the case of only one equation we saw that the two linearly independent fields were obtained as derivatives of one-parameter families of solutions from (2.16). However, as we are working with an ODE linear system with two equations, we expect to obtain four linearly independent Jacobi fields.

For the system, we can consider three natural one-parameter families of solutions. Indeed, similarly to (1.8), let

$$T \rightarrow v_\varepsilon(t + T)\Lambda \quad \text{and} \quad \varepsilon \rightarrow v_\varepsilon\Lambda.$$

The derivatives of these families provide us the solutions $\phi_{\varepsilon,0}^1 = \psi_{\varepsilon,0}^+\Lambda$ and $\phi_{\varepsilon,0}^2 = \psi_{\varepsilon,0}^-\Lambda$, where $\psi_{\varepsilon,0}^\pm$ are the previously described linearly independent Jacobi fields for one equation. Consequently our solutions are also linearly independent and have the same growth of the original fields.

Observe also that is possible to build another one-parameter family of solutions to (2.17). In fact, let $\Lambda(\alpha) = (\cos(\alpha + \alpha_0), \sin(\alpha + \alpha_0))$ with $\Lambda(0) = (\cos \alpha_0, \sin \alpha_0) = \Lambda$ be a path in the sphere and consider the family

$$\alpha \rightarrow v_\varepsilon \Lambda(\alpha).$$

If we differentiate this family with respect to the parameter we get a third the Jacobi field $\phi_{\varepsilon,0}^3 = v_\varepsilon \bar{\Lambda}$, where $\bar{\Lambda} = (-\Lambda_2, \Lambda_1)$. Note as the vectors Λ and $\bar{\Lambda}$ are orthogonal we automatically have that this field is linearly independent to the other two that we have already built. Moreover $\phi_{\varepsilon,0}^3$ is also a periodic field.

Unfortunately, we are only able to construct the last Jacobi field in an indirect way. Observe first that v_ε is a solution for the ODE $\bar{L}_{\varepsilon,0}v_\varepsilon = 0$, where

$$\bar{L}_{\varepsilon,0} = \partial_t^2 - \frac{(n-2)^2}{4} + \frac{n(n-2)}{4}v_\varepsilon^{\frac{4}{n-2}}. \quad (2.18)$$

We are writing the operator from Fowler equation in this manner intentionally. As this operator is linear we know that there exists a solution $\phi_{\varepsilon,0}^-$ to (2.18) which is linearly independent to $\phi_{\varepsilon,0}^+ := v_\varepsilon$. The last Jacobi field will be given by $\phi_0^4 = \phi_{\varepsilon,0}^-\bar{\Lambda}$. Also as $\bar{L}_{\varepsilon,0} < L_{\varepsilon,0}$, using standard methods of comparison theory for ODE's, we know that the growth of this field is at most linear.

Therefore we constructed four linearly independent Jacobi fields for the system with respect to the first eigenvalue.

Based on the construction above, the Jacobi fields for the higher eigenvalues λ_k will be given by $\phi_k^1 = \psi_{\varepsilon,k}^+\Lambda$, $\phi_k^2 = \psi_{\varepsilon,k}^-\Lambda$, where $\psi_{\varepsilon,k}^\pm$ are the original solutions constructed for one equation which we already know the growth. The remaining two solutions are $\phi_k^3 = \phi_{\varepsilon,k}^+\bar{\Lambda}$, $\phi_k^4 = \phi_{\varepsilon,k}^-\bar{\Lambda}$, where $\phi_{\varepsilon,k}^\pm$ are solutions of

$$\bar{L}_{\varepsilon,k}v = \partial_t^2v + \left(\frac{n(n-2)}{4}v_\varepsilon^{\frac{4}{n-2}} - \frac{(n-2)^2}{4} - \lambda_k \right) v = 0.$$

We know that the zero order term of the above operator is negative when $k > 0$ since $v_\varepsilon < 1$ and $\lambda_k \geq n-1$ for such k . This implies that $\bar{L}_{\varepsilon,k}$ satisfies the maximum principle, so we are able to determine the growth of these fields, concluding our analysis.

Chapter 3

Coupled elliptic system in the punctured ball

In this chapter we will study asymptotics of singular nonnegative solutions to the following system

$$\Delta u_i + \frac{n(n-2)}{4} |\mathcal{U}|^{\frac{4}{n-2}} u_i = 0 \quad (3.1)$$

in the punctured ball $B_1(0) \setminus \{0\}$, where $|\mathcal{U}|^2 = u_1^2 + u_2^2$ and Δ denotes the euclidian laplacian. This system is the natural generalization of the singular Yamabe problem in the flat case.

It will be convenient to consider v_i obtained from u_i by the change of variables of Fowler in order to work with the equivalent system

$$L_{cyl} v_i + \frac{n(n-2)}{4} |\mathcal{V}|^{\frac{4}{n-2}} v_i = 0, \quad (3.2)$$

where $L_{cyl} = \partial_t^2 + \Delta_{\mathbb{S}^{n-1}} - \frac{(n-2)^2}{4}$ is the cylindrical laplacian.

3.1 Upper bound

In this section our main purpose is to prove an upper bound for a solution \mathcal{U} of (3.1) which is singular at the origin. Such a result will be obtained as a direct consequence of an inequality for positive supersolutions of (3.1) that are smooth in an open set $\Omega \subset B_1(0)$. The complement $\Gamma = B_1(0) \setminus \Omega$ is then a relatively closed set in the unit ball.

Remember that a positive solution \mathcal{U} is a weak supersolution of (3.1) if it lies in $L^{(n+2)/(n-2)}(B_1(0))^2$ and if it satisfies

$$\Delta u_i + \frac{n(n-2)}{4} |\mathcal{U}|^{\frac{4}{n-2}} u_i \leq 0$$

in the distributional sense on $B_1(0)$.

We know that any solution \mathcal{U} which lies locally in $L^{\frac{n+2}{n-2}}(B_1(0))^2$ and tends uniformly to infinity near Γ may be extended to a weak supersolution in $B_1(0)$. Indeed, for any large constant L if we define \mathcal{U}_L whose coordinate functions are given by $u_{i,L} = \min\{u_i, L\}$ then

$$\Delta u_{i,L} \leq -\frac{n(n-2)}{4} |\mathcal{U}_L|^{\frac{4}{n-2}} u_{i,L} \chi_{|\mathcal{U}| \leq L},$$

where χ_E denotes the characteristic function of a set E . Applying the dominated convergence theorem we conclude then that \mathcal{U} defines a weak supersolution.

However, if we consider a solution \mathcal{U} of the system (3.1) which is smooth on an arbitrary open subset Ω , we can not conclude immediately that the solution lies locally in $L^{\frac{n+2}{n-2}}(B_1(0))^2$. In order to get this we will assume that the relatively closed set $\Gamma = B_1(0) \setminus \Omega$ is *thin*.

More precisely, let $\Gamma \subset B_1(0)$ be a relatively closed set of Lebesgue measure zero. We will say that $\Gamma \subset B_1(0)$ is a *thin* set if there exists a sequence ϕ_k of smooth functions in $B_1(0)$, such that

$$\begin{cases} \phi_k \equiv 0 & \text{in a neighborhood of } \Gamma \\ 0 \leq \phi_k \leq 1 \\ \phi_k \text{ converges to 1 in } B^n \setminus \Gamma \end{cases}$$

and

$$\lim_{k \rightarrow \infty} \int_{B_r(0)} \left(|\Delta \phi_k|^{\frac{n+2}{4}} + |\nabla \phi_k|^{\frac{n+2}{2}} \right) dx = 0.$$

for any $r < 1$. Observe that a submanifold of dimension at most $(n-2)/2$ is thin and consequently, a point is thin in any dimension $n \geq 2$.

Lemma 3.1 *Suppose \mathcal{U} is a solution of (3.1) on $\Omega = B_1(0) \setminus \Gamma$ where Γ is a relatively closed thin set. Then \mathcal{U} lies locally in $L^{\frac{n+2}{n-2}}(B_1(0))^2$, and defines a weak solution of (3.1) on $B_1(0)$.*

Proof. Let ξ be a nonnegative smooth function compactly supported in $B_1(0) \setminus \Gamma$. Multiplying the system (3.1) by $\xi^{\frac{n+2}{2}}$ and integrating by parts we have

$$-\int u_i \Delta(\xi^{\frac{n+2}{2}}) dx = \frac{n(n-2)}{4} \int |\mathcal{U}|^{\frac{4}{n-2}} u_i \xi^{\frac{n+2}{2}}$$

for each coordinate u_i , which taking the sum in i implies the bound

$$\int |\mathcal{U}|^{\frac{n+2}{n-2}} \xi^{\frac{n+2}{2}} dx \leq c \int \xi^{\frac{n-2}{2}} u(\xi |\Delta \xi| + |\nabla \xi|^2) dx$$

where $u = u_1 + u_2$. Applying Hölder's inequality we obtain

$$\int (\xi^{\frac{n-2}{2}} |\mathcal{U}|)^{\frac{n+2}{n-2}} dx \leq c \int (\xi |\Delta \xi| + |\nabla \xi|^2)^{\frac{n+2}{4}} dx. \quad (3.3)$$

Choose now $\xi = \psi \phi_k$, where ψ is a function which is equal to 1 in $B_r(0)$ for some $r < 1$ and equal to zero outside $B_{r_1}(0)$, for some $r < r_1$. Then the following estimate holds

$$(\xi |\Delta \xi| + |\nabla \xi|^2)^{\frac{n+2}{4}} \leq c(1 + |\Delta \phi_k|^{\frac{n+2}{4}} + |\nabla \phi_k|^{\frac{n+2}{2}}).$$

Replacing this estimate in (3.3), we have that

$$\int |\mathcal{U}|^{\frac{n+2}{n-2}} \xi^{\frac{n+2}{2}} dx \leq c \int (1 + |\Delta \phi_k|^{\frac{n+2}{4}} + |\nabla \phi_k|^{\frac{n+2}{2}}) dx \leq c$$

proving that each $u_i \in L^{\frac{n+2}{n-2}}(B_r(0))$ for any $r < 1$, and hence $u_i \in L^{\frac{n+2}{n-2}}_{loc}(B_1(0))$.

We will prove now that \mathcal{U} is a weak solution for the problem (3.1). In order to do this, given any smooth function ξ with compact support in B^n we need to show that

$$\int_{B_1(0)} \left(u_i \Delta(\xi) + \frac{n(n-2)}{4} \xi |\mathcal{U}|^{\frac{4}{n-2}} u_i \right) dx = 0.$$

Indeed, if we multiply the system (3.1) by $\xi \phi_k$ and integrate by parts, we obtain

$$\int_{B_1(0)} \left(u_i \Delta(\xi \phi_k) + \frac{n(n-2)}{4} \xi \phi_k |\mathcal{U}|^{\frac{4}{n-2}} u_i \right) dx = 0.$$

We may calculate

$$\Delta(\xi \phi_k) = \xi \Delta \phi_k + 2 \langle \nabla \xi, \nabla \phi_k \rangle + \phi_k \Delta \xi.$$

By Hölder's inequality it follows that

$$\left| \int u_i \xi \Delta(\phi_k) dx \right| \leq \|u_i \xi\|_{L^{(n+2)/(n-2)}} \|\Delta(\phi_k)\|_{L^{(n+2)/4}(\text{supp } \xi)},$$

and the right-hand side tends to zero as k tends to infinity. Similarly

$$\left| \int u_i \langle \nabla \xi, \nabla \phi_k \rangle dx \right| \leq \|u_i\| \|\nabla \xi\|_{L^{(n+2)/(n-2)}} \|\nabla \phi_k\|_{L^{(n+2)/4}(\text{supp} \xi)}$$

and the right-hand side converges to zero. Therefore applying the dominated convergence theorem and letting $k \rightarrow \infty$ we conclude the result. ■

Another key ingredient in order to obtain the upper bound is the existence of spherical solutions on \mathbb{R}^n . Remember that for any $\mu > 0$ and $x_0 \in \mathbb{R}^n$, the *spherical solution* of dilatation μ and center x_0 is given by

$$u_{\mu, x_0}(x) = \left(\frac{2\mu}{1 + \mu^2|x - x_0|^2} \right)^{\frac{n-2}{2}}. \quad (3.4)$$

As mentioned in the introduction of this thesis, O. Druet, E. Hebey and J. Vetois [7] proved that positive solutions of (3.1) in \mathbb{R}^n are given by a spherical solution times a vector Λ in the unit sphere with positive coordinates. We can now state and prove the upper bound estimate.

Theorem 3.2 *Let \mathcal{U} be a positive supersolution of (3.1) which is a smooth solution on an open set $\Omega \subset B_1(0)$. Then there exists a positive constant c such that*

$$|\mathcal{U}|(x) \leq cd(x, \Gamma)^{\frac{2-n}{2}} \left(\inf_{\partial B_{3/4}(0)} |\mathcal{U}| \right)^{-1}$$

for all $x \in \Omega \cap B_{1/2}(0)$.

Proof. Denote by Γ_ε the neighborhood with radius ε about Γ . For $x \in B_{5/8}(0)$, set $U_\varepsilon(x) = d_\varepsilon(x)^{\frac{n-2}{2}} |\mathcal{U}|(x)$ where $d_\varepsilon(x) = \min\{d(x, \Gamma_\varepsilon), \frac{5}{8} - |x|\}$. In order to prove the Theorem it is enough to show that

$$\sup_{B_{5/8}(0)} U_\varepsilon \leq c \left(\inf_{\partial B_{3/4}(0)} |\mathcal{U}| \right)^{-1} \quad (3.5)$$

for some constant c not depending on ε .

Let $\eta = |\mathcal{U}(x_0)|^{-\frac{2}{n-2}}$, where x_0 is a maximum point of U_ε and consider the rescaled functions

$$w_{i,\eta}(x) = \eta^{\frac{n-2}{2}} u_i(x_0 + \eta x).$$

The first step to prove our result is the following claim.

Claim 1: Let $R > 0$ and $\delta \in (0, \frac{1}{2}]$ be constants. There exists a constant c_1 depending only of n , R and δ such that if $M_\varepsilon \equiv \sup_{B(0, \frac{5}{8})} U_\varepsilon(x) \geq c_1$ then

$$\|u_{\mu, y_0} \Lambda_i - w_{i, \eta}\|_{C^2(B_R(0))} < \delta, \quad (3.6)$$

where u_{μ, y_0} is the spherical solution defined in (3.4) with $|y_0| \leq c_2$ and $1/c_2 \leq \mu \leq c_2$, and $\Lambda = (\Lambda_1, \Lambda_2)$ is a vector in the unit sphere with nonnegative coordinates.

Suppose by contradiction that this claim is not true. Then there would exist a sequence of maximum points $x_k \in B_{5/8}(0)$ of U_ε such that $U_\varepsilon(x_k)$ converges to infinity and (3.6) does not hold for any $\mu > 0$ and any $y_0 \in \mathbb{R}^n$. Let

$$w_{i, k}(x) = \eta_k^{\frac{n-2}{2}} u_i(x_k + \eta_k x)$$

for each coordinate, where $\eta_k = |\mathcal{U}(x_k)|^{-\frac{2}{n-2}}$. Since x_k is a maximum point if $r_k = \frac{1}{2}d_\varepsilon(x_k)$, we have that

$$|\mathcal{U}(x)| \leq 2^{\frac{n-2}{2}} |\mathcal{U}(x_k)|,$$

for $x \in B_{r_k}(x_k)$. Since $w_{i, k}$ is a solution of (3.1) which is regular and bounded, standard elliptic theory then implies that, after passing to a subsequence, $w_{i, k}$ converge in C^2 norm on compact subsets of \mathbb{R}^n to a positive solution $u_{i, 0}$ to

$$\Delta u_{i, 0} + \frac{n(n-2)}{4} |\mathcal{U}_0|^{\frac{4}{n-2}} u_{i, 0} = 0,$$

which by a theorem due to Druet and Hebey [6] is the spherical standard solution.

Now let us prove that the bounds on y_0 and μ are valid. Indeed, from the previous argument we know that

$$|\mathcal{W}_\lambda| \leq 2^{\frac{n-2}{2}} \text{ in } B_{R'}(0),$$

where $R' = d_\varepsilon(x_0)/2\eta \geq c_1^{\frac{2}{n-2}}/2$. We can always assume that $c_1 \geq 2^{n-2}$ and consequently $R' \geq 2$. Then, it holds

$$|\mathcal{U}_{\mu, y_0}(y)| \leq 2^{\frac{n-2}{2}} + \delta \leq 2^{\frac{n}{2}} \quad (3.7)$$

for all $y \in B_2(0)$. On the other hand, we have

$$|\mathcal{U}_{\mu, y_0}(0)| \geq 1 - \delta \geq 1/2. \quad (3.8)$$

Note that the lower bound $2^{-\frac{2}{n-2}} \leq 2\mu$ follows directly from (3.8). In order to prove the upper bound suppose first that $y_0 \in B_2(0)$ then, taking $y = y_0$ in (3.7), we

get that $2\mu \leq 2^{\frac{n}{n-2}}$. Otherwise, there would exist $y_1 \in B_2(0)$ with $|y_1 - y_0|^2 = |y_0|^2 - 1$ and taking $y = y_1$ in (3.7) we have

$$2^{-\frac{2}{n-2}}\mu \leq 1 + \mu^2|y_0|^2 - \mu^2 \leq 2^{\frac{n}{n-2}}\mu - \mu^2.$$

This implies that $\mu \leq 2^{\frac{n}{n-2}}$ and thus $|y_0| \leq 2^{n/(n-2)}$. This finishes the proof of the claim.

Observe now that, if we shift the center point x_0 to a new point x_1 which is within distance $2c_2\eta$, the inequality (3.6) can be replaced by

$$\|u_{1,0}\Lambda_i - w_\lambda\|_{C^2(B_R(0))} < \delta. \quad (3.9)$$

Indeed, $|\mathcal{U}_{\mu,y_0}|$ has a nondegenerate maximum point at y_0 . Then we conclude that for k sufficiently large there exists a nondegenerate maximum point y_k of $|\mathcal{W}_k|$ near y_0 . Since y_0 and μ are bounded there will be a corresponding local maximum point x_1 of $|\mathcal{W}_k|$ satisfying $|x_0 - x_1| \leq 2c_2\eta < 1/16$, increasing c_1 if necessary. If we redefine the functions $w_{i,k}$ replacing x_0 by x_1 , the result follows.

In order to prove (3.5), observe first that since \mathcal{U} is a supersolution, $|\mathcal{U}|$ is locally bounded in $L^{(n+2)/(n-2)}(B^n)$, such function is superharmonic and thus it can be redefined on Γ as to be upper semicontinuous, which implies that $\inf_{\partial B_{3/4}(0)} |\mathcal{U}|$ is bounded from above.

Let $R = e^2$ and $\delta \in (0, \frac{1}{2}]$ be fixed constants. If the left-hand side of inequality (3.5) is bounded by c_1 (given in the Claim 1), then the result follows directly. Suppose that this is not the case.

From now on we will work in cylindrical coordinates. If $\eta = |\mathcal{U}(x_1)|^{-\frac{2}{n-2}}$ then we have

$$v_i(t, \theta) = |x|^{\frac{n-2}{2}} w_{i,\eta}(x),$$

where $t = -\log|x|$ and $\theta = x/|x|$.

Since x_1 lies in the ball of radius $11/16$ about the origin, its distance to $\partial B_{3/4}(0)$ is at least $1/16$. Then the ball $B_{\eta^{-1}/16}(0)$ is contained in the domain of $w_{i,\eta}$. Now define,

$$\mathcal{V}_{1,0}(t) = |x|^{\frac{n-2}{2}} u_{1,0}(x) \Lambda = (\cosh t)^{\frac{2-n}{2}} \Lambda.$$

From (3.9), we know that \mathcal{V} is close in the C^2 norm on $[-2, \infty) \times \mathbb{S}^{n-1}$ to the function $\mathcal{V}_{1,0}$. Consequently $\partial_t v_i(-1, \theta) > 0$ for all $\theta \in \mathbb{S}^{n-1}$.

We will apply the Alexandrov technique to the coordinates v_i on the region $[-\log(\frac{1}{16}\eta^{-1}), \infty) \times \mathbb{S}^{n-1}$, reflecting across the spheres $\{\alpha\} \times \mathbb{S}^{n-1}$ starting with α sufficiently large, and continuing as far as possible. Once $\partial_t v_i(-1, \theta) > 0$, this procedure must end before α reaches -1. Let

$$v = v_1 + v_2,$$

$$v_{i,\alpha}(t, \theta) = v_i(2\alpha - t, \theta)$$

and

$$v_\alpha(t, \theta) = v(2\alpha - t, \theta).$$

Then, the difference between the function v_i and the reflection satisfies

$$L(v_i - v_{i,\alpha}) + \frac{n(n-2)}{4} |\mathcal{V}|^{\frac{4}{n-2}} v_i - \frac{n(n-2)}{4} |\mathcal{V}_\alpha|^{\frac{4}{n-2}} v_{i,\alpha} = 0. \quad (3.10)$$

Note that

$$|\mathcal{V}|^{\frac{4}{n-2}} v - |\mathcal{V}_\alpha|^{\frac{4}{n-2}} v_\alpha = \sum_{j=1}^2 b_{i,\alpha}(v_i - v_{i,\alpha}),$$

where

$$b_{i,\alpha} = v(v_i + v_{i,\alpha}) \frac{|\mathcal{V}|^{\frac{4}{n-2}} - |\mathcal{V}_\alpha|^{\frac{4}{n-2}}}{|\mathcal{V}|^2 - |\mathcal{V}_\alpha|^2} + |\mathcal{V}_\alpha|^{\frac{4}{n-2}} > 0.$$

By (3.10) we have

$$L(v - v_\alpha) + \sum_{i=1}^2 b_{i,\alpha}(v_i - v_{i,\alpha}) = 0$$

weakly. In order to conclude the proof, we will also need the following:

Claim 2: There exist $\alpha_0 > -1$ and $\theta_0 \in \mathbb{S}^{n-1}$ such that

$$v(-\log(\frac{1}{16}\eta^{-1}), \theta_0) = v_{\alpha_0}(-\log(\frac{1}{16}\eta^{-1}), \theta_0).$$

Indeed, define

$$\alpha_0 = \inf\{\alpha_1; (v - v_\alpha)(t, \theta) \geq 0 \text{ in } [-\log(\frac{1}{16}\eta^{-1}), \alpha_1], \forall \alpha \geq \alpha_1\}.$$

By continuity, $(v - v_{\alpha_0})(t, \theta) \geq 0$ in $[-\log(\frac{1}{16}\eta^{-1}), \alpha_0]$. If we suppose by contradiction that the claim is not true, then

$$(v - v_{\alpha_0})(-\log(\frac{1}{16}\eta^{-1}), \theta) > 0$$

for all $\theta \in \mathbb{S}^{n-1}$. Since the following inequality

$$L(v - v_{\alpha_0}) - D(v - v_{\alpha_0}) \leq - \sum_{i=1}^2 b_{i,\alpha_0}(v_i - v_{i,\alpha_0}) \leq 0, \quad (3.11)$$

holds weakly, where D is a positive function, this implies that $(v - v_{\alpha_0})(t, \theta) > 0$ for every $-\log((16\eta)^{-1}) < t < \alpha_0$ and $\theta \in \mathbb{S}^{n-1}$, since on the boundary of $[-\log((16\eta)^{-1}), \alpha_0]$ the difference is nonnegative.

From the definition of α_0 , there exist a sequence $\{\alpha_j\}$ such that $\alpha_j < \alpha_0$ and $\alpha_j \rightarrow \alpha_0$, and a sequence $\{(t_j, \alpha_j)\}$ of interior minimum points of $v - v_{\alpha_j}$ such that $(t_j, \theta_j) \rightarrow (t^*, \theta^*)$ with $(v - v_{\alpha_j})(t_j, \theta_j) < 0$. Taking the limit $j \rightarrow \infty$ we get $(v - v_{\alpha_0})(t^*, \theta^*) = 0$ and $\nabla(v - v_{\alpha_0})(t^*, \theta^*) = 0$. Therefore $t^* = \alpha_0$, which is a contradiction to Hopf's Lemma. This proves the Claim 2.

Hence we see that there exist α_0 and θ_0 such that

$$v(-\log(\frac{1}{16}\eta^{-1}), \theta_0) = v_{\alpha_0}(-\log(\frac{1}{16}\eta^{-1}), \theta_0),$$

which implies

$$\inf \left\{ v(-\log(\frac{1}{16}\eta^{-1}), \theta) : \theta \in \mathbb{S}^{n-1} \right\} \leq \sup \left\{ v(t, \theta) : t > \log(\frac{1}{16}\eta^{-1}) - 2\alpha_0, \theta \in \mathbb{S}^{n-1} \right\}. \quad (3.12)$$

Note that since \mathcal{U} is superharmonic, it holds

$$\inf \left\{ v(-\log(\frac{1}{16}\eta^{-1}), \theta) : \theta \in \mathbb{S}^{n-1} \right\} = 16^{\frac{2-n}{2}} \inf_{\partial B_{1/16}(x_1)} u \geq 16^{\frac{2-n}{2}} \inf_{\partial B_{3/4}(0)} |\mathcal{U}|.$$

On the other hand, for η small and fixed, we have $|\mathcal{V}|(t, \theta) \leq 2|\mathcal{V}_{1,0}|(t, \theta)$ for $t \geq 0$, and therefore

$$\inf_{\partial B_{3/4}(0)} |\mathcal{U}| \leq c \sup_{[\log(\frac{1}{16}\eta^{-1}) - 2\alpha_0, \infty)} |\mathcal{V}_{1,0}| \leq c\eta^{\frac{n-2}{2}} |\mathcal{V}_{1,0}| = c|\mathcal{U}(x_1)|^{-1}.$$

Since U_ε is bounded by a constant times $|\mathcal{U}|$, the inequality (3.5) is true, which finishes the proof of the result.

■

The following result follows directly as a consequence of Lemma 3.1 and Theorem 3.2.

Corollary 3.3 *Assume that \mathcal{U} is a nonnegative smooth solution of (3.1) defined in the punctured unit ball $B_1^n(0) \setminus \{0\}$. Either \mathcal{U} extends as a smooth solution to the ball, or there exists positive constants $c > 0$, such that*

$$|\mathcal{U}|(x) \leq c|x|^{\frac{2-n}{2}},$$

for $0 < |x| < 1/2$.

Corollary 3.4 *Suppose \mathcal{U} is a positive smooth solution of (3.1) in $\Omega = B_1^n(0) \setminus \{0\}$. Then there exists a constant $c > 0$ such that*

$$\max_{|x|=r} u_i \leq c \min_{|x|=r} u_i \quad (3.13)$$

for every $0 < r < \frac{1}{4}$. Moreover, $|\nabla u_i| \leq c|x|^{-1}u_i$ and $|\nabla^2 u_i| \leq c|x|^{-2}u_i$.

Proof. Define $u_{i,r}(y) = r^{\frac{n-2}{2}} u_i(ry)$, for every $0 < r < \frac{1}{4}$ and $|y| < r^{-1}$. Then the upper bound given by previous theorem implies that $u_{i,r}(y) \leq c|y|^{\frac{2-n}{2}}$, for $|y| < \frac{1}{2}r^{-1}$. In particular, if $\frac{1}{2} \leq |y| \leq \frac{3}{2}$, we have that $u_{i,r}(y) \leq 2^{\frac{n-2}{2}}c$. Moreover

$$\Delta u_{i,r}(y) + \frac{n(n-2)}{4} |\mathcal{U}_r(y)|^{\frac{4}{n-2}} u_{i,r}(y) = 0.$$

Hence the Harnack inequality for linear elliptic equations and standard elliptic theory imply there exists $c > 0$, not depending on r , such that

$$\max_{|x|=1} u_{i,r} \leq c \min_{|x|=1} u_{i,r},$$

and $|\nabla u_{i,r}| + |\nabla^2 u_{i,r}| \leq cu_{1,r}$ on the sphere of radius 1. This finishes the proof of the corollary. ■

3.2 Pohozaev invariant and lower bound

In this section we will prove a lower bound for the solutions of (3.1) in the punctured ball and we will give some information about the Pohozaev invariant when the singularity is nonremovable.

We will begin by proving a Lemma that gives us a sufficient condition so that the origin can be a removable singularity.

Lemma 3.5 *Let \mathcal{U} be a positive solution of (3.1) defined in the punctured ball and denote $u(x) = u_1(x) + u_2(x)$. If*

$$\lim_{|x| \rightarrow 0} |x|^{\frac{n-2}{2}} u(x) = 0, \quad (3.14)$$

then \mathcal{U} extends as a smooth solution to all of B_1 .

Proof. Observe first that Lemma 3.1 implies that \mathcal{U} extends as a weak solution on the unit ball. By hypothesis (3.14) we know that $v(t, \theta)$ converges uniformly to zero as t tends to ∞ , where v corresponds to u by the change of variables of Fowler (1.2). Since v satisfies

$$\partial_t^2 v + \Delta_{\mathbb{S}^{n-1}} v - \frac{(n-2)^2}{4} v + \frac{n(n-2)}{4} |\mathcal{V}|^{\frac{4}{n-2}} v = 0,$$

there exist $\beta > 0$ and $t_0 > 0$ such that

$$Lv = \partial_t^2 v + \Delta_{\mathbb{S}^{n-1}} v \geq \beta v$$

for $t \geq t_0$. Consider now the function

$$w = ce^{-\sqrt{\beta}t} + \varepsilon e^{\sqrt{\beta}t},$$

where c is chosen such that $ce^{-\sqrt{\beta}t_0} > v(t_0, \theta)$ for all $\theta \in \mathbb{S}^{n-1}$. Note that w satisfies $Lw = \beta w$. Since w is growing at an exponential rate and v is converging to zero, we can choose sufficiently large T such that $v(T, \theta) < w(T, \theta)$ for all θ in the unit sphere.

Applying the maximum principle we get for all $t_0 \leq t \leq T$ and for all θ that

$$v(t, \theta) \leq ce^{-\sqrt{\beta}t} + \varepsilon e^{\sqrt{\beta}t}. \quad (3.15)$$

Since T was chosen arbitrarily large, this inequality holds for all $t \geq t_0$. Making ε converge to zero, we conclude that for all $t \geq t_0$

$$e^{\sqrt{\beta}t} v(t, \theta) \leq c,$$

which implies $u(x) \leq c|x|^q$ for $q = (2-n)/2 + \sqrt{\beta}$. This estimate shows that $u \in L_{loc}^p(B_1(0))$ for arbitrarily large p and then by elliptic theory the function u extends smoothly across the origin. Consequently, each coordinate function u_i is smooth. ■

Similarly to what we did in the previous chapter we can define for $r > 0$ the Pohozaev integral as

$$P(r, \mathcal{U}) = \int_{\partial B_r} \left(\frac{n-2}{2} \left\langle \mathcal{U}, \frac{\partial \mathcal{U}}{\partial \nu} \right\rangle - \frac{r}{2} |\nabla \mathcal{U}|^2 + r \left| \frac{\partial \mathcal{U}}{\partial \nu} \right|^2 + r \frac{(n-2)^2}{8} |\mathcal{U}|^{\frac{2n}{n-2}} \right) d\sigma \quad (3.16)$$

where ν is the unit outer normal of ∂B_r . Multiplying each equation of (3.1) by $x \cdot \nabla u_i$ and integrating over $B_r \setminus B_s$ we verify for all $r, s > 0$ that

$$P(r, \mathcal{U}) = P(s, \mathcal{U})$$

and hence we can denote it by $P(\mathcal{U})$ and call it the *Pohozaev invariant*.

Theorem 3.6 *Assume that \mathcal{U} is a nonnegative smooth solution of (3.1) defined in the punctured unit ball $B_1^n(0) \setminus \{0\}$. Either \mathcal{U} extends as a smooth solution to the ball, or there exists a positive constant c such that*

$$|\mathcal{U}|(x) \geq c|x|^{\frac{2-n}{2}}.$$

In the latter case the radial Pohozaev invariant of \mathcal{U} is negative.

Proof. Again it will be more convinient work in the cylindrical background. If $v = v_1 + v_2$, define

$$\bar{v}(t) = \int_{\mathbb{S}^{n-1}} v(t, \theta) d\theta.$$

As a consequence of the spherical Harnack inequality given by Corollary 3.4, we know that v/\bar{v} is bounded from above. Suppose that the solution \mathcal{U} can not be extended smoothly across the origin. By Lemma 3.5, this imply that either the claimed lower bound holds, or there exists a sequence t_k of critical points of v such that $\lim_{k \rightarrow \infty} \bar{v}(t_k) = 0$.

On one hand, in the cylindrical setting the Pohozaev integral $P(t_k, \mathcal{V}) = P(e^{-t_k}, \mathcal{U})$ is given by

$$P(t_k, \mathcal{V}) := \int_{t_k \times \mathbb{S}^{n-1}} \left(\frac{1}{2} |\partial_t \mathcal{V}|^2 - \frac{1}{2} |\nabla_\theta \mathcal{V}|^2 - \frac{(n-2)^2}{8} |\mathcal{V}|^2 + \frac{(n-2)^2}{8} |\mathcal{V}|^{\frac{2n}{n-2}} \right) d\sigma.$$

Consequently using the fact that the Pohozaev integral is independent of k , the Harnack inequality, standard elliptic estimates as well as the fact that v_i is a solution of (3.2) imply that the invariant must vanish.

On the other hand, consider the sequence of functions

$$w_{i,k}(t, \theta) = \frac{v_i(t + t_k, \theta)}{\bar{v}(t_k)}.$$

Using elliptic estimates, we can extract a subsequence of $w_{i,k}$ which converges in C^2 topology on compact subsets to a positive solution of $L_{cyl} w_i = 0$. In euclidian coordinates the associated function $h_i(x)$ defined on $\mathbb{R}^n \setminus \{0\}$ is then a positive harmonic function, and can therefore be written as $h_i(x) = a_i |x|^{2-n} + b_i$, for some $a_i, b_i \geq 0$. That is,

$$w_i(t, \theta) = a_i e^{\frac{n-2}{2}t} + b_i e^{-\frac{n-2}{2}t}.$$

We also know from the fact that the sequence $\{t_k\}$ was chosen as critical points for \bar{v} that $a_1 + a_2 = b_1 + b_2$ is positive. Then, by a direct computation

$$\begin{aligned} \lim_{k \rightarrow \infty} \int_{\mathbb{S}^{n-1}} \left[\sum_i \left(\frac{1}{2} (\partial_t w_{i,k})^2 - \frac{1}{2} |\nabla_\theta w_{i,k}|^2 - \frac{(n-2)^2}{8} w_{i,k}^2 \right) + \bar{v}(t_k)^{\frac{4}{n-2}} |\mathcal{W}_k|^{\frac{2n}{n-2}} \right] d\theta \\ = \int_{\mathbb{S}^{n-1}} \sum_i \left(\frac{1}{2} (\partial_t w_i)^2 - \frac{(n-2)^2}{8} w_i^2 \right) d\theta = -\omega_n \frac{(n-2)^2}{2} \langle a, b \rangle. \end{aligned}$$

We claim that $\langle a, b \rangle$ is positive. Indeed, if we consider the function

$$z_{i,k}(x) = |x|^{\frac{2-n}{2}} w_{i,k}(-\log|x|, x/|x|)$$

one verifies that $\mathcal{Z}_k = (z_{1,k}, z_{2,k})$ satisfies the following system

$$\Delta z_{i,k} + \frac{n(n-2)}{4} |\mathcal{Z}_k|^{\frac{4}{n-2}} z_{i,k} = 0 \quad \text{in} \quad B_{r_k^{-1}}(0) \setminus \{0\}, \quad (3.17)$$

where $r_k = e^{-t_k}$.

Now, multiplying line i of system (3.17) by $z_{j,k}$ and integrating over $B_1(0) \setminus B_\varepsilon(0)$, we obtain that

$$\int_{B_1 \setminus B_\varepsilon} z_{j,k} \Delta z_{i,k} dx = \frac{n(n-2)}{4} \int_{B_1 \setminus B_\varepsilon} |\mathcal{Z}_k|^{\frac{4}{n-2}} z_{i,k} z_{j,k} dx.$$

Reversing i and j and subtracting one equation from the other give us

$$\int_{B_1 \setminus B_\varepsilon} (z_{j,k} \Delta z_{i,k} - z_{i,k} \Delta z_{j,k}) dx = 0,$$

which, integrating by parts, guarantees that

$$\int_{\partial B_1} (z_{j,k} \partial_r z_{i,k} - z_{i,k} \partial_r z_{j,k}) dx = \int_{\partial B_\varepsilon} (z_{j,k} \partial_r z_{i,k} - z_{i,k} \partial_r z_{j,k}) dx. \quad (3.18)$$

In order to analyse the last integral on the right-hand side, let

$$\varphi_{i,k}^\varepsilon(y) = \varepsilon^{\frac{n-2}{2}} z_{i,k}(\varepsilon y).$$

It holds

$$\int_{\partial B_\varepsilon} (z_{j,k} \partial_r z_{i,k} - z_{i,k} \partial_r z_{j,k}) dx = \int_{\partial B_1} (\varphi_{j,k}^\varepsilon \partial_r \varphi_{i,k}^\varepsilon - \varphi_{i,k}^\varepsilon \partial_r \varphi_{j,k}^\varepsilon) dx.$$

Since

$$\Delta \varphi_{i,k}^\varepsilon + \frac{n(n-2)}{4} |\varphi_k^\varepsilon|^{\frac{4}{n-2}} \varphi_{i,k}^\varepsilon = 0$$

in $B_{(\varepsilon r_k)^{-1}}(0) \setminus \{0\}$, by Corollary 3.3 we get

$$|\varphi_k^\varepsilon(y)| \leq C|y|^{\frac{2-n}{2}}.$$

By standard elliptic estimates, after passing to a subsequence, $\varphi_{i,k}^\varepsilon$ converges in C^2 topology locally in compact subsets of $\mathbb{R}^n \setminus \{0\}$, to a positive solution

$$\Delta u_i + \frac{n(n-2)}{4} |\mathcal{U}|^{\frac{4}{n-2}} u_i = 0$$

which using Theorem 2.10 as well as [7, Proposition 1.1] is of the form $u_0 \Lambda$, where $\Lambda = (\Lambda_1, \Lambda_2)$ is a vector in the unit sphere with nonnegative coordinates. Hence

$$\lim_{\varepsilon \rightarrow 0} \int_{\partial B_1} (\varphi_{j,k}^\varepsilon \partial_r \varphi_{i,k}^\varepsilon - \varphi_{i,k}^\varepsilon \partial_r \varphi_{j,k}^\varepsilon) d\sigma_g = \int_{\partial B_1} (u_0 \partial_r u_0 \Lambda_i \Lambda_j - u_0 \partial_r u_0 \Lambda_j \Lambda_i) d\sigma_g = 0.$$

Making $\varepsilon \rightarrow 0$ in (3.18), we conclude that

$$\int_{\partial B_1} (z_{j,k} \partial_r z_{i,k} - z_{i,k} \partial_r z_{j,k}) dx = 0. \quad (3.19)$$

Finally, we conclude that

$$\int_{\partial B_1} (h_j \partial_r h_i - h_i \partial_r h_j) d\sigma_g = 0,$$

and consequently

$$a_1 b_2 = a_2 b_1$$

which finishes the proof of our claim.

Note now that this is a contradiction, for this is also a limit of rescalings of the Pohozaev invariant for \mathcal{V} , and hence must be zero. Indeed

$$0 = \bar{v}(t_k)^{-2} P(\mathcal{V}) = \int_{\mathbb{S}^{n-1}} \left[\sum_i \left(\frac{1}{2} (\partial_t w_{i,k})^2 - \frac{1}{2} |\nabla_\theta w_{i,k}|^2 - \frac{(n-2)^2}{8} w_{i,k}^2 \right) + \bar{v}(t_k)^{\frac{4}{n-2}} |\mathcal{W}_k|^{\frac{2n}{n-2}} \right] d\theta \neq 0,$$

for sufficiently large k . This establishes the lower bound.

In order to show that the Pohozaev invariant is negative, by the upper and lower bounds, we can choose a sequence \tilde{t}_k tending to ∞ so that the corresponding translated solutions $t \rightarrow \mathcal{V}(t + t_k, \theta)$ converge in C^2 norm on compact subsets of $\mathbb{R} \times \mathbb{S}^{n-1}$ to a Fowler-type solution \mathcal{V}_0 satisfying the same bounds and defined on all of $\mathbb{R} \times \mathbb{S}^{n-1}$. Therefore

$$P(\mathcal{V}) = P(\mathcal{V}_0) < 0,$$

which finishes the proof. \blacksquare

3.3 Simple convergence to a radial solution

Our main goal in this section is to prove that a singular solution to the system (3.1) is asymptotic to a radial Fowler-type solution near the nonremovable isolated singularity.

Theorem 3.7 *Suppose that \mathcal{U} is a solution of the system (3.1) in the punctured ball $B_1^n(0) \setminus \{0\}$. If there exist positive constants c_1 and c_2 such that*

$$c_1|x|^{\frac{2-n}{2}} \leq |\mathcal{U}|(x) \leq c_2|x|^{\frac{2-n}{2}} \quad (3.20)$$

then there exist a Fowler-type solution $\mathcal{U}_0 = u_0\Lambda$ of (6), where u_0 is a Fowler solution such that

$$\mathcal{U}(x) = (1 + O(|x|^\alpha))\mathcal{U}_0(x) \quad (3.21)$$

as $x \rightarrow 0$, for some $\alpha > 0$.

Proof. First we observe that (3.20) implies that the origin is a nonremovable singularity. Thus, we get that the Pohozaev invariant is negative.

Consider v_i obtained from u_i by the change of variables from Fowler and let $\{\tau_k\}$ be a sequence of real numbers such that $\tau_k \rightarrow \infty$. Consider the translated sequence $v_{i,k}(t, \theta) = v_i(t + \tau_k, \theta)$ defined in $(-\tau_k, \infty) \times \mathbb{S}^{n-1}$. By (3.20), we get that

$$c_1 \leq |\mathcal{V}_k(t, \theta)| \leq c_2,$$

where $\mathcal{V}_k = (v_{1,k}, v_{2,k})$. Consequently, by standard elliptic estimates, we get the uniform boundedness of any derivative for $t > 0$ and moreover there exists a subsequence, also denoted by $v_{i,k}$, which converges in the C_{loc}^2 topology, to a positive solution of

$$L_{cyl}v_{i,0} + \frac{n(n-2)}{4}|\mathcal{V}_0|^{\frac{4}{n-2}}v_{i,0} = 0$$

defined in the whole cylinder. By the characterization result given by Theorem 2.10, such limit is a Fowler-type solution and we know that there exists a Fowler solution v_ε and a vector in the unit sphere with positive coordinates Λ such that $\mathcal{V}_\varepsilon(t) = \Lambda v_\varepsilon(t)$. Hence \mathcal{V}_ε does not depend on θ , and we necessarily have that any angular derivative $\partial_\theta v_{i,k}$ converges uniformly to zero.

Besides, we claim that

$$\begin{aligned} v_{i,k}(t, \theta) &= \bar{v}_{i,k}(t)(1 + o(1)) \\ \nabla v_{i,k}(t, \theta) &= -\bar{v}'_{i,k}(t)(1 + o(1)), \end{aligned}$$

as $t \rightarrow \infty$. In fact, suppose that the first equality above is false. Then there exist $\bar{\varepsilon} > 0$ and sequences $\tau_k \rightarrow \infty$, $\theta_k \rightarrow \theta \in \mathbb{S}^{n-1}$ such that

$$\left| \frac{v_{i,k}(\tau_k, \theta_k)}{\bar{v}_{i,k}(\tau_k)} - 1 \right| \geq \bar{\varepsilon}$$

for some $i \geq 1$. This is a contradiction because, after passing to a subsequence, \mathcal{V}_k converges to a rotationally symmetric Fowler-type solution \mathcal{V}_0 . The second inequality follows from similar arguments.

Note now that

$$P(\mathcal{V}_\varepsilon) := P(0, \mathcal{V}_\varepsilon) = \lim_{k \rightarrow \infty} P(0, \mathcal{V}_k) = \lim_{k \rightarrow \infty} P(\tau_k, \mathcal{V}) = P(\mathcal{V}). \quad (3.22)$$

So we can conclude that the necksize ε of the limit function is independent of the sequence of numbers τ_k . Therefore, for each sequence $\tau_k \rightarrow \infty$ the correspondent sequence \mathcal{V}_k converges to a function $\mathcal{V}_{\varepsilon,T}(t) = \Lambda v_\varepsilon(t+T)$, with $\Lambda \in \mathbb{S}_+^1$, for some $T \in \mathbb{R}$ which depends on the sequence τ_k .

We will show that there exists $T_0 \in \mathbb{R}$ such that \mathcal{V}_k converges to $\mathcal{V}_{\varepsilon,T_0}$ for any sequence $\tau_k \rightarrow \infty$. The idea is to use a delicate rescaling argument due originally to Leon Simon. In order to do that we will prove several claims using the Jacobi fields studied in section 2.5 as a tool.

Let T_ε be the period of \mathcal{V}_ε and $A_\tau = \sup_{t \geq 0} |\partial_\theta \mathcal{V}_\tau|$, where $\mathcal{V}_\tau(t, \theta) = \mathcal{V}(t+\tau, \theta)$. Note that $A_\tau < \infty$, since $|\partial_\theta \mathcal{V}_\tau|$ converges uniformly to zero as $t \rightarrow \infty$.

Claim 1: For every $c > 0$, there exists a positive integer N such that, for any $\tau > 0$, either

(i) $A_\tau \leq ce^{-2\tau}$ or

(ii) A_τ is attained at some point in $I_N \times \mathbb{S}^{n-1}$, where $I_N = [0, NT_\varepsilon]$.

Suppose the Claim is not true. Then there exist a constant $c > 0$ and sequences $\tau_k, s_k \rightarrow \infty, \theta_k \in \mathbb{S}^{n-1}$ such that $|\partial_\theta \mathcal{V}_\tau|(s_k, \theta_k) = A_{\tau_k}$ and $A_{\tau_k} > ce^{-2\tau_k}$ as $k \rightarrow \infty$. Then we can translate back further s_k and define $\tilde{v}_{i,k}(t, \theta) = v_{i,k}(t + s_k, \theta)$. Define $\varphi_{i,k} = A_{\tau_k}^{-1} \partial_\theta \tilde{v}_{i,k}$ and note that $|\varphi_{\mathbf{k}}| \leq 1$, where $\varphi_{\mathbf{k}} = (\varphi_{1,k}, \varphi_{2,k})$. Now, we have

$$L_{cyl}(\tilde{v}_{i,k}) + \frac{n(n-2)}{4} |\tilde{\mathcal{V}}_k|^{\frac{4}{n-2}} \tilde{v}_{i,k} = 0,$$

where the quantities with tilde are the originals replacing t by $t + \tau_k + s_k$. Taking the derivative with respect to θ and multiplying by $A_{\tau_k}^{-1}$, we get

$$L_{cyl}(\varphi_{i,k}) + \frac{n(n-2)}{4} |\tilde{\mathcal{V}}_k|^{\frac{4}{n-2}} \varphi_{i,k} + n |\tilde{\mathcal{V}}_k|^{\frac{4}{n-2}-2} \tilde{v}_{ik} \langle \tilde{\mathcal{V}}_k, \boldsymbol{\varphi}_k \rangle = 0$$

Now we can use elliptic theory to extract a subsequence $\varphi_{i,k}$ which converges in compact subsets to a nontrivial and bounded Jacobi field $\boldsymbol{\varphi} = (\varphi_1, \varphi_2)$ which satisfies the following system

$$L_{cyl}(\varphi_i) + \frac{n(n-2)}{4} v_\varepsilon^{\frac{4}{n-2}} \varphi_i + n \Lambda_i \langle \Lambda, \boldsymbol{\varphi} \rangle v_\varepsilon^{\frac{4}{n-2}} = 0.$$

Since each coordinate function of the limit φ_i has no zero eigencomponent relative to Δ_θ , we get a contradiction because a Jacobi field with such property is necessarily unbounded. This proves the Claim 1.

Now suppose we have a sequence $v_{i,k}(t, \theta)$ converging to $\Lambda_i v_\varepsilon(t + T)$ as $k \rightarrow \infty$. Define

$$w_{i,k}(t, \theta) = v_{i,k}(t, \theta) - \Lambda_i v_\varepsilon(t + T)$$

and set

$$\eta_k = b \max_{I_N} |(w_{1,k}, w_{2,k})| \quad \text{and} \quad \varphi_{i,k} = \eta_k^{-1} w_{i,k},$$

where $b > 0$ is a fixed number to be chosen later and so that $|(\varphi_{1,k}, \varphi_{2,k})| \leq b^{-1}$ on I_N . Then

$$L_{cyl}(w_{i,k}) + \frac{n(n-2)}{4} \left(|\mathcal{V}_k|^{\frac{4}{n-2}} v_{i,k} - \Lambda_i v_\varepsilon^{\frac{n+2}{n-2}} \right) = 0 \quad (3.23)$$

and

$$|\mathcal{V}_k|^{\frac{4}{n-2}} v_{i,k} - \Lambda_i v_\varepsilon^{\frac{n+2}{n-2}} = |\mathcal{V}_k|^{\frac{4}{n-2}} w_{i,k} + \Lambda_i v_\varepsilon \frac{|\mathcal{V}_k|^{\frac{4}{n-2}} - v_\varepsilon^{\frac{4}{n-2}}}{|\mathcal{V}_k|^2 - v_\varepsilon^2} \sum_j w_{j,k} (v_{j,k} + \Lambda_i v_\varepsilon).$$

Multiplying (3.23) by η_k^{-1} and taking the limit $k \rightarrow \infty$ we get

$$L_{cyl}(\varphi_i) + \frac{n(n-2)}{4} v_0^{\frac{4}{n-2}} \varphi_i + n \Lambda_i \langle \Lambda, \boldsymbol{\varphi} \rangle v_\varepsilon^{\frac{4}{n-2}} = 0,$$

on the whole cylinder, where $\boldsymbol{\varphi} = (\varphi_1, \varphi_2)$ is a Jacobi field.

Claim 2: The Jacobi field $\boldsymbol{\varphi}$ is bounded for $t \geq 0$.

To prove this claim we will use the analysis done in section 2.5. By the spectral decomposition for the laplacian in the sphere, we know that it is possible to write the Jacobi field as

$$\boldsymbol{\varphi} = a_1 \phi_{\varepsilon,0}^1 + a_2 \phi_{\varepsilon,0}^2 + a_3 \phi_{\varepsilon,0}^3 + a_4 \phi_{\varepsilon,0}^4 + \tilde{\boldsymbol{\varphi}}$$

where $\phi_{\varepsilon,0}^i$ are the linearly independent Jacobi fields corresponding to the eigencomponent independent of θ , and $\tilde{\varphi}$ denotes the projection onto the orthogonal complement. We also know that the functions $\phi_{\varepsilon,0}^1$ and $\phi_{\varepsilon,0}^3$ are bounded and $\phi_{\varepsilon,0}^2$ and $\phi_{\varepsilon,0}^4$ are linearly growing.

Let us show that $\tilde{\varphi}$ is bounded by proving that each $\partial_\theta \tilde{\varphi}_i = \partial_\theta \varphi_i$ is bounded for $t \geq 0$. In fact, the function $\partial_\theta \varphi_i$ is the limit of $\eta_k^{-1} \partial_\theta v_{i,k}$, and we can suppose that $\partial_\theta \varphi_i$ is nontrivial, otherwise the result is immediate.

If the first item of Claim 1 happens then

$$\sup_{t \geq 0} (\eta_k^{-1} |\partial_\theta v_{i,k}|) \leq c \eta_k^{-1} e^{-2\tau_k} \leq C.$$

While if the second item of Claim 1 is true then

$$\sup_{t \geq 0} (\eta_k^{-1} |\partial_\theta v_{i,k}|) \leq \sup_{t \geq 0} (\eta_k^{-1} |\partial_\theta \mathcal{V}_k|) = \sup_{t \in I_N} (\eta_k^{-1} |\partial_\theta \mathcal{V}_k|) \leq C,$$

since the sequence $\eta_k^{-1} |\partial_\theta \mathcal{V}_k|$ converges in the C_{loc}^2 topology. Therefore each $\tilde{\varphi}_i$ is bounded for $t \geq 0$, hence exponentially decaying.

To finish the proof of the Claim 2 we need to show that $a_2 = a_4 = 0$. To see this note that the convergence $\varphi_{i,k} = \eta_k^{-1} w_{i,k} \rightarrow \varphi_i$ implies that

$$\begin{aligned} \mathcal{V}_k &= \Lambda v_{\varepsilon,T} + \eta_k \varphi + o(\eta_k) \\ &= \Lambda v_{\varepsilon,T} + \eta_k (a_1 \phi_{\varepsilon,0}^1 + a_2 \phi_{\varepsilon,0}^2 + a_3 \phi_{\varepsilon,0}^3 + a_4 \phi_{\varepsilon,0}^4 + \tilde{\varphi}) + o(\eta_k), \end{aligned}$$

where $v_{\varepsilon,T}(t) = v_\varepsilon(t + T)$. On the other hand we have that

$$P(0, \mathcal{V}_k) = P(\tau_k, \mathcal{V}) = P(\mathcal{V}) + O(e^{-2\tau_k}) = P(v_{\varepsilon,T}) + O(e^{-2\tau_k}).$$

Since $\lim_{k \rightarrow \infty} (\eta_k^{-1} e^{-2\tau_k}) = 0$, we would have a contradiction in case a_2 or a_4 is not zero because the two sides of the last equality would differ for sufficiently large k . Thus each φ is bounded for $t \geq 0$.

Now we show that there exists some T so that the difference between \mathcal{V} and $\mathcal{V}_{\varepsilon,T} = \Lambda v_{\varepsilon,T}$ goes to zero as $t \rightarrow \infty$. Since we do not know the correct translation parameter, define $\mathcal{V}_\tau(t, \theta) = \mathcal{V}(t + \tau, \theta)$ and $\mathcal{W}_\tau(t, \theta) = \mathcal{V}_\tau(t, \theta) - \Lambda v_\varepsilon(t)$. Let $C_1 > 0$ be a fixed constant and consider the interval I_N as in the Claim 1. Set also $\eta(\tau) = b \max_{I_N} |\mathcal{W}_\tau|$, where $b > 0$ is again a positive constant to be chosen later. We observe that $\eta(\tau) \rightarrow 0$ as $\tau \rightarrow \infty$. Let us prove the following claim.

Claim 3: If N , b and τ are sufficiently large and η is sufficiently small, then there exists s with $|s| \leq C_1\eta(\tau)$ so that $\eta(\tau + NT_\varepsilon + s) \leq \frac{1}{2}\eta(\tau)$.

Suppose the claim is not true. Then there exists some sequence $\tau_k \rightarrow \infty$ such that $\eta(\tau_k) \rightarrow 0$ and for any s satisfying $|s| \leq C_1\eta(\tau_k)$ we have that $\eta(\tau_k + NT_\varepsilon + s) > \frac{1}{2}\eta(\tau_k)$. Define $\varphi_{i,k} = \eta(\tau_k)^{-1}w_{i,\tau_k}$, similarly to the previous claim. We can suppose that $\varphi_{i,k}$ converges in C^∞ on compact sets to a Jacobi field, which by Claim 2 is bounded for $t \geq 0$. So we can write

$$\varphi = a_1\phi_{\varepsilon,0}^1 + a_3\phi_{\varepsilon,0}^3 + \tilde{\varphi} \quad (3.24)$$

where $\tilde{\varphi}$ has exponential decay. Note that $|\varphi| \leq b^{-1}$ on I_N , which implies that a_1 and a_3 are uniformly bounded, independently of the sequence τ_k . Moreover, since $\phi_{\varepsilon,0}^1 = v'_\varepsilon\Lambda$ and $\phi_{\varepsilon,0}^3 = v_\varepsilon\bar{\Lambda}$ we have

$$|a_3v_\varepsilon| \leq |\langle \varphi, \bar{\Lambda} \rangle| + |\langle \tilde{\varphi}, \bar{\Lambda} \rangle| \leq b^{-1} + |\tilde{\varphi}| \quad (3.25)$$

on I_N . We know that $v_\varepsilon \geq \varepsilon$ and $\tilde{\varphi}$ decreases exponentially with a fixed rate, and so we can choose b and N sufficiently large such that $|a_3|$ is sufficiently small.

Set $s_k = -\eta(\tau_k)a_1$ whose absolute value is less than $C_1\eta(\tau_k)$ if we choose C_1 sufficiently large. Hence for $t \in [0, 2NT_\varepsilon]$ we have

$$\begin{aligned} \mathcal{W}_{\tau_k+s_k}(t, \theta) &= \mathcal{V}(t + \tau_k - \eta(\tau_k)a_1, \theta) - \Lambda v_\varepsilon(t) \\ &= \mathcal{V}_{\tau_k}(t - \eta(\tau_k)a_1, \theta) - \Lambda v_\varepsilon(t - \eta(\tau_k)a_1) \\ &\quad - \eta(\tau_k)a_1 \Lambda \frac{v_\varepsilon(t - \eta(\tau_k)a_1) - v_\varepsilon(t)}{-\eta(\tau_k)a_1} \\ &= \eta(\tau_k)\varphi_k(t - \eta(\tau_k)a_1, \theta) - \eta(\tau_k)a_1\phi_{\varepsilon,0}^1 + o(\eta(\tau_k)) \\ &= \mathcal{W}_{\tau_k}(t, \theta) - \eta(\tau_k)a_1\phi_{\varepsilon,0}^1 + o(\eta(\tau_k)), \end{aligned}$$

where $\varphi_k = (\varphi_{1,k}, \varphi_{2,k})$. Here we used the equality $\mathcal{W}_{\tau_k} = \eta(\tau_k)\varphi_k + o(\eta(\tau_k))$ and the fact that $\varphi_k(t - \eta(\tau_k)a_1, \theta) - \varphi_k(t, \theta)$ goes to zero as $\tau_k \rightarrow \infty$.

Consequently, by (3.24), for $t \in [0, 2NT_\varepsilon]$ we get that

$$\mathcal{W}_{\tau_k+s_k} = \eta(\tau_k)\tilde{\varphi} + \eta(\tau_k)a_3\phi_{\varepsilon,0}^3 + o(\eta(\tau_k)),$$

which implies

$$\max_{I_N} |\mathcal{W}_{\tau_k+s_k+NT_\varepsilon}| = \max_{[NT_\varepsilon, 2NT_\varepsilon]} |\mathcal{W}_{\tau_k+s_k}| \leq \eta(\tau_k) \max_{[NT_\varepsilon, 2NT_\varepsilon]} (|\tilde{\varphi}| + |a_3v_\varepsilon|) + o(\eta(\tau_k)).$$

Since $\tilde{\varphi}$ decreases exponentially with a fixed rate, by (3.25) we can choose N and $b > 0$ sufficiently large in a way that the last equality implies that

$$\max_{I_N} |\mathcal{W}_{\tau_k + s_k + NT_\varepsilon}| \leq \frac{1}{4} \eta(\tau_k).$$

which on its turn gives $\eta(\tau + NT_\varepsilon + s) \leq \frac{1}{2} \eta(\tau)$, a contradiction. This ends the proof of the Claim 3.

Once the claim above is proved, using an iterative argument, we are ready to prove that there exists σ such that $w_{i,\sigma} \rightarrow 0$ as $t \rightarrow \infty$ for each coordinate. First choose τ_0 and N sufficiently large satisfying the Claim 3 and such that $C_1 \eta(\tau_0) \leq \frac{1}{2} NT_\varepsilon$. Let $s_0 = -\eta(\tau_0) a_1$ be chosen as above. Thus we have $|s_0| \leq C_1 \eta(\tau_0) \leq \frac{1}{2} NT_\varepsilon$. Define inductively three sequences by

$$\begin{aligned} \sigma_k &= \tau_0 + \sum_{i=0}^{k-1} s_i \\ \tau_k &= \tau_{k-1} + s_{k-1} + NT_\varepsilon = \sigma_k + kNT_\varepsilon \\ s_k &= -\eta(\tau_k) a_1. \end{aligned}$$

By the Claim 3 we get by induction $\eta(\tau_k) \leq 2^{-k} \eta(\tau_0)$ and $|s_k| \leq 2^{-k-1} NT_\varepsilon$. Hence there exists the limit $\sigma = \lim \sigma_k \leq \tau_0 + NT_\varepsilon$ and then $\tau_k \rightarrow \infty$ as $k \rightarrow \infty$.

We claim σ is the correct translation parameter. In fact, choose k such that $t = kNT_\varepsilon + [t]$ with $[t] \in I_N$, and write

$$\begin{aligned} w_{i,\sigma}(t, \theta) &= v_i(t + \sigma, \theta) - \Lambda_i v_\varepsilon(t) \\ &= v_i(t + \sigma, \theta) - v_i(t + \sigma_k, \theta) + v_i(t + \sigma_k, \theta) - \Lambda_i v_\varepsilon(t). \end{aligned}$$

Since $\partial_t v_i$ is uniformly bounded, we have

$$v_i(t + \sigma, \theta) - v_i(t + \sigma_k, \theta) = \partial_t v_i(t_0) \sum_{j=k}^{\infty} s_j = O(2^{-k}),$$

for some t_0 . Besides,

$$v_i(t + \sigma_k, \theta) - \Lambda_i v_\varepsilon(t) = v_i(\tau_k + [t], \theta) - \Lambda_i v_\varepsilon([t]) = w_{i,\tau_k}([t], \theta).$$

Thus,

$$\mathcal{W}_\sigma(t, \theta) = \mathcal{W}_{\tau_k}([t], \theta) + O(2^{-k}).$$

Since $b \max_{I_N} |\mathcal{W}_{\tau_k}| = \eta(\tau_k) \leq \eta(\tau_k) \leq 2^{-k} \eta(\tau_0)$, it holds $|w_{i,\sigma}(t, \theta)| = O(2^{-k})$ or equivalently, using that $t = kNT_\varepsilon + [t]$, we have

$$|w_{i,\sigma}(t, \theta)| \leq C_1 e^{-\frac{\log 2}{NT_\varepsilon} t}$$

which finishes the proof of the theorem. ■

Chapter 4

Yamabe-type system in the punctured ball

Let g be a smooth Riemannian metric on the unit ball $B_1^n(0) \subset \mathbb{R}^n, n \geq 3$. Consider a positive solution to the system

$$\Delta_g u_i - \sum_{j=1}^2 A_{ij}(x) u_j + \frac{n(n-2)}{4} |\mathcal{U}|^{\frac{4}{n-2}} u_i = 0 \quad (4.1)$$

in the punctured ball $\Omega = B_1^n(0) \setminus \{0\}$ and $\mathcal{U} = (u_1, u_2)$. In order to prove our main result, during the proof of the results we will need sometimes that the potential A satisfy some of the following conditions.

(H1) $-A$ is *cooperative*, that is, the components in the nondiagonal A_{ij} of A , $i \neq j$, are nonpositive;

(H2) In dimension $n = 5$, there exists a C^2 -function f such that

$$A(x) = f(x) Id_2 + O(|x|) \quad (4.2)$$

near the origin, where Id_2 is the identity matrix.

4.1 Upper bound near a singularity

In this section we will obtain upper and lower bounds for solutions to our system defined in the punctured ball. Considering the diffeomorphism $\Phi(t, \theta) = e^{-t}\theta$ between

the half cylinder and the punctured ball introduced in the first chapter, define the change of variables from Fowler as

$$v_i(t, \theta) = |x|^{\frac{n-2}{2}} u_i(x)$$

and $\hat{g} = e^{2t}\Phi^*g = (e^{\frac{n-2}{2}t})^{\frac{4}{n-2}}\Phi^*g$. Using that

$$L_{v^{\frac{4}{n-2}}g}(u) = v^{-\frac{n+2}{n-2}}L_g(vu), \quad (4.3)$$

where the linear operator $L_g = \Delta_g - \frac{n-2}{4(n-1)}R_g$ is the *conformal laplacian*, we obtain that the system is equivalent to

$$\mathcal{L}_{\hat{g}}(v_i) - \sum_{j=1}^2 B_{ij}v_j + \frac{n(n-2)}{4}|\mathcal{V}|^{\frac{4}{n-2}}v_i = 0, \quad (4.4)$$

where

$$\begin{aligned} \mathcal{L}_{\hat{g}}(v_i) &= \Delta_{\hat{g}}v_i - \frac{n-2}{4(n-1)}(R_{\hat{g}} - e^{-2t}R_{\Phi^*g})v_i, \\ B_{ij} &= e^{-2t}A_{ij} \circ \Phi \end{aligned} \quad (4.5)$$

and $\mathcal{V} = (v_1, v_2)$.

It is also useful to remember that in cylindrical coordinates we have

$$R_{\hat{g}} = (n-2)(n-1) + 2(n-1)e^{-t}\frac{\partial_r\sqrt{|g|}}{\sqrt{|g|}} + e^{-2t}R_g \circ \Phi,$$

and so

$$R_{\hat{g}} - e^{-2t}R_{\Phi^*g} = (n-2)(n-1) + 2(n-1)e^{-t}\frac{\partial_r\sqrt{|g|}}{\sqrt{|g|}} \circ \Phi. \quad (4.6)$$

Theorem 4.1 *Suppose $3 \leq n \leq 5$. Assume that $\mathcal{U} = (u_1, u_2)$ is a positive solution of (1) in $\Omega = B_1^n(0) \setminus \{0\}$. There exists a constant $c > 0$ such that*

$$|\mathcal{U}|(x) \leq cd_g(x, 0)^{\frac{2-n}{2}}, \quad (4.7)$$

for $0 < d_g(x, 0) < \frac{1}{2}$.

Proof. Given $x_0 \in \Omega$ with $d_g(x_0, 0) < \frac{1}{2}$ and $s \in (0, \frac{1}{4})$ such that $\bar{B}_s(x_0) \subset \Omega$, define

$$f(x) = (s - d_g(x, x_0))^{\frac{n-2}{2}}|\mathcal{U}(x)|,$$

for $x \in \bar{B}_s(x_0)$. It suffices to show that there exists a positive constant C such that any such f satisfies $f(x) \leq C$ in $B_s(x_0)$. To see this, taking $s = \frac{|x_0|}{2}$ gives $f(x_0) = s^{\frac{n-2}{2}}|\mathcal{U}(x_0)| \leq C$.

The proof will be by contradiction, so assume there is no such constant C . Then we can find a sequence of points $x_{0,k}$ and positive numbers s_k so that, if $x_{1,k}$ denotes the maximum point of the corresponding f_k , we have

$$f_k(x_{1,k}) = (s_k - d_g(x, x_{0,k}))^{\frac{n-2}{2}} |\mathcal{U}(x_{1,k})| \rightarrow \infty.$$

Note that, $0 < s_k < \frac{1}{4}$ implies that $(s_k - d_g(x, x_{0,k}))^{\frac{n-2}{2}} < 2^{2-n}$ and therefore $2^{n-2} f_k(x) < |\mathcal{U}(x)|$. Hence $|\mathcal{U}(x_{1,k})| \rightarrow \infty$ and consequently $x_{1,k} \rightarrow 0$.

Let $\varepsilon_k = |\mathcal{U}(x_{1,k})|^{\frac{-2}{n-2}}$ and define,

$$\tilde{u}_{i,k}(y) = \varepsilon_k^{\frac{n-2}{2}} u_i(\exp_{x_{1,k}}(\varepsilon_k y)).$$

for all $i \in \{1, 2\}$. Note that $|\tilde{\mathcal{U}}_k(0)| = 1$, where $\tilde{\mathcal{U}}_k = (\tilde{u}_{1,k}, \tilde{u}_{2,k})$. Also note that the function $\tilde{u}_{i,k}$ is defined for all y such that $|y| \leq \varepsilon_k^{-1}(s_k - d_g(x_{0,k}, x_{1,k}))$. Moreover, if

$$d_g(x, x_{1,k}) \leq r_k = \frac{1}{2}(s_k - d_g(x_{0,k}, x_{1,k}))$$

then

$$d_g(x, x_{0,k}) - d_g(x_{0,k}, x_{1,k}) \leq d_g(x, x_{1,k}) \leq r_k = \frac{1}{2}(s_k - d_g(x_{0,k}, x_{1,k})),$$

and this implies that

$$d_g(x, x_{0,k}) \leq s_k + \frac{1}{2}(-s_k + d_g(x_{0,k}, x_{1,k})).$$

Thus

$$r_k \leq s_k - d_g(x_{0,k}, x_{1,k}).$$

Therefore

$$r_k^{\frac{n-2}{2}} |\mathcal{U}(x)| \leq f_k(x) \leq f_k(x_{1,k}) = (2r_k \varepsilon_k^{-1})^{\frac{n-2}{2}} = (2r_k)^{\frac{n-2}{2}} |\mathcal{U}(x_{1,k})| \rightarrow \infty,$$

and

$$|\mathcal{U}(x)| \leq 2^{\frac{n-2}{2}} |\mathcal{U}(x_{1,k})|$$

for all x with $d_g(x, x_{1,k}) \leq r_k$. It follows that

$$\tilde{u}_{i,k}(y) = |\mathcal{U}(x_{1,k})|^{-1} u_i(\exp_{x_{1,k}}(\varepsilon_k y)) \leq |\mathcal{U}(x_{1,k})|^{-1} |\mathcal{U}(x)| \leq 2^{\frac{n-2}{2}},$$

for all y with $|\varepsilon_k y| \leq r_k$, that is, $|y| \leq r_k \varepsilon_k^{-1} \rightarrow \infty$.

Now if we define $(\tilde{g}_k)_{lm}(y) := g_{lm}(\varepsilon_k y)$, then $\tilde{u}_{i,k}$ satisfies

$$\Delta_{\tilde{g}_k} \tilde{u}_{i,k}(y) - \varepsilon_k^2 \sum_{j=1}^2 \tilde{A}_{ij}^k(y) \tilde{u}_{j,k}(y) + \frac{n(n-2)}{4} |\tilde{\mathcal{U}}_k(y)|^{\frac{4}{n-2}} \tilde{u}_{i,k}(y) = 0, \quad (4.8)$$

where $\tilde{A}_{ij}^k(y) := A_{ij}(\exp_{x_{1,k}}(\varepsilon_k y))$.

Standard elliptic theory then implies that, after passing to a subsequence, $\{\tilde{u}_{i,k}\}_k$ converge in C^2 norm on compact subsets of \mathbb{R}^n to a positive solution $u_{i,0}$ to

$$-\Delta u_{i,0} = \frac{n(n-2)}{4} |\mathcal{U}_0|^{\frac{4}{n-2}} u_{i,0},$$

which satisfies $|\mathcal{U}_0(0)| = 1$ and $u_{i,0}(y) \leq 2^{\frac{n-2}{2}}$ for every $y \in \mathbb{R}^n$. By a theorem due to Druet and Hebey [6], we can conclude that there exist $a \in \mathbb{R}^n$, $\mu > 0$ and $\Lambda \in \mathbb{S}_+^1$ such that

$$\mathcal{U}_0(y) = \left(\frac{2\mu}{1 + \mu^2|y - a|^2} \right)^{\frac{n-2}{2}} \Lambda.$$

Since $|\mathcal{U}_0(0)| = 1$, we conclude that $|a| \leq 1$ and $\mu \in [1/2, 1]$.

Now note that \mathcal{U}_0 has a nondegenerate maximum point at a . Then we conclude that there is a sequence $y_k \rightarrow a$ such that y_k is a nondegenerate maximum point of $|\tilde{\mathcal{U}}_k|$. We can assume $|y_k| \leq 2$ and therefore there will be a corresponding local maximum point $x_{2,k}$ of $|\mathcal{U}|$ satisfying $d_g(x_{2,k}, x_{1,k}) \leq 2\varepsilon_k$. If we redefine the functions $\tilde{u}_{i,k}$ replacing $x_{1,k}$ by $x_{2,k}$ we get as before that a subsequence $\{\tilde{u}_{i,k}\}_k$ converge in the C^2 norm on compact subsets of \mathbb{R}^n to

$$\mathcal{U}_0(y) = \left(\frac{1}{1 + \frac{1}{4}|y|^2} \right)^{\frac{n-2}{2}} \Lambda.$$

Note that, by construction, we have that $|x_{2,k}| < 7/8$, so we can consider $\tilde{u}_{i,k}$ as defined for $|y| \leq \frac{1}{16}\varepsilon_k^{-1}$, with a possible singularity at some point on the sphere of radius $|x_{2,k}|\varepsilon_k^{-1} \rightarrow \infty$, where now $\varepsilon_k = |\mathcal{U}(x_{2,k})|^{-\frac{2}{n-2}}$.

Let us introduce

$$v_{i,k}(t, \theta) = |y|^{\frac{n-2}{2}} \tilde{u}_{i,k}(y),$$

where $t = -\log|y|$ and $\theta = \frac{y}{|y|}$. These functions are defined for $t > \log(\frac{1}{16}\varepsilon_k^{-1})$, with a singularity at some point (t'_k, θ'_k) , $t'_k = \log(|x_{2,k}\varepsilon_k^{-1}|)$. Now define,

$$\mathcal{V}_0(t) = |y|^{\frac{n-2}{2}} \mathcal{U}_0(y) = \left(e^t + \frac{1}{4}e^{-t} \right)^{\frac{2-n}{2}} \Lambda.$$

Since $\tilde{\mathcal{U}}_k \rightarrow \tilde{\mathcal{U}}_0$ in the C_{loc}^2 topology, we know that given $R > 0$ the inequalities

$$\begin{aligned} |\mathcal{V}_k(t, \theta) - \mathcal{V}_0(t)| &\leq R^{-1} e^{\frac{2-n}{2}t}, \\ |\partial_t \mathcal{V}_k(t, \theta) - \mathcal{V}'_0(t)| &\leq R^{-1} e^{\frac{2-n}{2}t}, \\ |\partial_t^2 \mathcal{V}_k(t, \theta) - \mathcal{V}''_0(t)| &\leq R^{-1} e^{\frac{2-n}{2}t}, \\ |\partial_{\theta_i} \mathcal{V}_k(t, \theta)| &\leq R^{-1} e^{\frac{2-n}{2}t}, \\ |\partial_{\theta_i \theta_m}^2 \mathcal{V}_k(t, \theta)| &\leq R^{-1} e^{\frac{2-n}{2}t} \end{aligned}$$

are satisfied for $t > -\log R$ and sufficiently large k .

In particular

$$\partial_t v_{i,k}(-\log 3n, \theta) = \frac{2-n}{2} \left((3n)^{-1} + \frac{3n}{4} \right)^{\frac{2-n}{2}} - R^{-1} (3n)^{\frac{2-n}{2}} > 0 \quad (4.9)$$

for all $\theta \in \mathbb{S}^{n-1}$ and for $R > 0$ large enough.

For a fixed small number $\delta > 0$ to be chosen later, define

$$\Gamma_k = [\log(\delta \varepsilon_k^{-1}), \infty) \times \mathbb{S}^{n-1}.$$

Since $\tilde{\mathcal{U}}_k \rightarrow \mathcal{U}_0$ in the C_{loc}^2 topology and

$$\mathcal{V}_0(\log(\delta \varepsilon_k^{-1})) = \left(\delta^{-1} \varepsilon_k + \frac{\delta \varepsilon_k^{-1}}{4} \right)^{\frac{2-n}{2}} \Lambda \geq c(\delta) > 0,$$

we obtain

$$\mathcal{V}_k(\log(\delta \varepsilon_k^{-1})) \geq c(\delta) > 0.$$

We will apply the Alexandrov technique to $v_{i,k}$ on the region Γ_k reflecting across the spheres $\{\lambda\} \times S^{n-1}$. To simplify the notation we will drop the subscript k .

Define $\Gamma_\lambda = [-\log(\delta \varepsilon^{-1}), \lambda]$, \hat{g}_λ the pull-back of the metric \hat{g} by the reflection across the sphere $\{\lambda\} \times S^{n-1}$

$$\begin{aligned} v &= v_1 + v_2 \\ v_{i,\lambda}(t, \theta) &= v_i(2\lambda - t, \theta) \end{aligned}$$

and

$$v_\lambda(t) = v(2\lambda - t, \theta).$$

Remember that $\hat{g} = e^{2t} \Phi^* \tilde{g}$, and $(\tilde{g}_k)_{lm}(y) = g_{lm}(\varepsilon_k y)$.

By (4.4) we can write

$$\begin{aligned} \mathcal{L}_{\hat{g}}(v_i - v_{i,\lambda}) - \sum_{j=1}^2 B_{ij}(v_j - v_{j,\lambda}) + \frac{n(n-2)}{4} |\mathcal{V}|^{\frac{4}{n-2}} v_i - \frac{n(n-2)}{4} |\mathcal{V}_\lambda|^{\frac{4}{n-2}} v_{i,\lambda} \\ = (\mathcal{L}_{\hat{g}_\lambda} - \mathcal{L}_{\hat{g}})v_{i,\lambda} - \sum_{j=1}^2 (B_{ij}^\lambda - B_{ij})v_{j,\lambda}. \end{aligned} \quad (4.10)$$

Note that,

$$|\mathcal{V}|^{\frac{4}{n-2}} v - |\mathcal{V}_\lambda|^{\frac{4}{n-2}} v_\lambda = \sum_{j=1}^2 b_{i,\lambda}(v_i - v_{i,\lambda}),$$

where

$$b_{i,\lambda} = v(v_i + v_{i,\lambda}) \frac{|\mathcal{V}|^{\frac{4}{n-2}} - |\mathcal{V}_\lambda|^{\frac{4}{n-2}}}{|\mathcal{V}|^2 - |\mathcal{V}_\lambda|^2} + |\mathcal{V}_\lambda|^{\frac{4}{n-2}} > 0.$$

By (4.10) we have

$$\mathbb{L}_{\hat{g}}(v_1 - v_{1,\lambda}, v_2 - v_{2,\lambda}) + \sum_{i=1}^2 b_{i,\lambda}(v_i - v_{i,\lambda}) = Q_\lambda,$$

where

$$Q_\lambda = (\mathcal{L}_{\hat{g}_\lambda} - \mathcal{L}_{\hat{g}})v_\lambda + \sum_{i,j=1}^2 (B_{ij}^\lambda - B_{ij})v_{j,\lambda} \quad (4.11)$$

and

$$\mathbb{L}_{\hat{g}}(w_1, w_2) = \mathcal{L}_{\hat{g}} \left(\sum_{j=1}^2 w_j \right) - \sum_{i,j=1}^2 B_{ij} w_j. \quad (4.12)$$

Claim 1: There exists a constant $c_1 > 0$, not depending on δ , such that $|Q_\lambda(t, \theta)| \leq q_\lambda(t) = c_1 \varepsilon^2 e^{\frac{n-6}{2}t} e^{(2-n)\lambda}$.

First, we observe that $\frac{\partial_r \sqrt{|g|}}{\sqrt{|g|}} = O(|x|)$ and $v_\lambda(t) = O(e^{\frac{2-n}{2}(2\lambda-t)})$ and this implies that

$$\begin{aligned} |R_{\hat{g}_\lambda} - e^{-2t} R_{\Phi^* \tilde{g}_\lambda} - (R_{\hat{g}} - e^{-2t} R_{\Phi^* \tilde{g}})| v_\lambda(t, \theta) &\leq C \varepsilon^2 e^{-2t} e^{\frac{2-n}{2}(2\lambda-t)} \\ &= C \varepsilon^2 e^{\frac{n-6}{2}t} e^{(2-n)\lambda}. \end{aligned}$$

On the other hand, since $\hat{g} = e^{2t} \Phi^* \tilde{g}$ and $\tilde{g}_{ij} = \delta_{ij} + O(\varepsilon^2 |y|^2)$ in normal coordinates, we have $\hat{g} = dt^2 + d\theta^2 + O(\varepsilon^2 e^{-2t})$. It follows that

$$|(\Delta_{\hat{g}_\lambda} - \Delta_{\hat{g}})v_\lambda| \leq C \varepsilon^2 e^{\frac{n-6}{2}t} e^{(2-n)\lambda}.$$

Also by (4.5) we have

$$\left| \sum_{i,j=1}^2 (B_{ij}^\lambda - B_{ij})v_{j,\lambda} \right| \leq C \varepsilon^2 e^{\frac{n-6}{2}t} e^{(2-n)\lambda},$$

proving the first claim.

Claim 2: Suppose $3 \leq n \leq 5$, and let $\gamma > 0$ be a small number. Then there exists a family of functions $h_\lambda(t)$, defined on Γ_λ , satisfying the following properties

$$h_\lambda(\lambda) = 0; \quad (4.13)$$

$$h_\lambda \geq 0; \quad (4.14)$$

$$\mathbb{L}_{\hat{g}}(h_\lambda, h_\lambda) \geq Q_\lambda; \quad (4.15)$$

$$h_\lambda \leq v_i - v_{i,\lambda}, \text{ if } \lambda \text{ is sufficiently large} \quad (4.16)$$

and

$$h_\lambda(-\log(\delta\varepsilon^{-1})) \leq c_3 \max \left\{ \varepsilon^{\frac{n-2}{2}}, \varepsilon^{\frac{6-n}{2}-\gamma} \right\}, \quad (4.17)$$

for some positive constant c which depends only on δ .

We will begin the proof of the second claim with some useful computations for the operator $\mathbb{L}_{\hat{g}}$ in order to get the desired estimates. Remember that $\hat{g} = e^{\frac{2-n}{2}t} \Phi^* \tilde{g}$, thus by (4.3) we obtain

$$L_{\hat{g}}f = e^{-\frac{n+2}{2}t} L_{\Phi^* \tilde{g}}(e^{\frac{n-2}{2}t} f) = e^{-\frac{n+2}{2}t} \left(\Delta_{\tilde{g}}(|y|^{\frac{2-n}{2}} \tilde{f}) - \frac{n-2}{4(n-1)} |y|^{\frac{2-n}{2}} R_{\tilde{g}} \tilde{f} \right) \circ \Phi,$$

where $\tilde{f} = f \circ \Phi^{-1}$. Hence

$$\mathcal{L}_{\hat{g}}f = L_{\hat{g}}f + \frac{n-2}{4(n-1)} e^{-2t} R_{\Phi^*g}f = e^{-\frac{n+2}{2}t} \Delta_{\tilde{g}}(|y|^{\frac{2-n}{2}} \tilde{f}) \circ \Phi.$$

So, if f depends only on t , then we can use the expression of the laplacian for radial functions given by

$$\Delta_{\tilde{g}}u = \partial_r^2 u + \frac{n-1}{r} \partial_r + \frac{\partial_r \sqrt{|\tilde{g}|}}{\sqrt{|\tilde{g}|}} \partial_r u.$$

Note that

$$\partial_r(|y|^{\frac{2-n}{2}} \tilde{f}) = \frac{2-n}{2} |y|^{-\frac{n}{2}} \tilde{f} + |y|^{\frac{2-n}{2}} \partial_r \tilde{f}$$

and

$$\partial_r^2(|y|^{\frac{2-n}{2}} \tilde{f}) = \frac{n(n-2)}{4} |y|^{-\frac{n+2}{2}} \tilde{f} + (2-n) |y|^{-\frac{n}{2}} \partial_r \tilde{f} + |y|^{\frac{2-n}{2}} \partial_r^2 \tilde{f}.$$

On the other hand, since $\tilde{f}(|y|) = f(-\log |y|)$, we have

$$\partial_r \tilde{f} = -|y|^{-1} f' \circ \Phi^{-1}$$

and

$$\partial_r^2 \tilde{f} = |y|^{-2} f' \circ \Phi^{-1} + |y|^{-2} f'' \circ \Phi^{-1}.$$

Then we get

$$\partial_r(|y|^{\frac{2-n}{2}} \tilde{f}) = \frac{2-n}{2} |y|^{-\frac{n}{2}} f \circ \Phi^{-1} - |y|^{\frac{n}{2}} f' \circ \Phi^{-1}$$

and

$$\partial_r^2(|y|^{\frac{n-2}{2}} \tilde{f}) = \frac{n(n-2)}{4} |y|^{-\frac{n+2}{2}} f \circ \Phi^{-1} + (n-1) |y|^{-\frac{n+2}{2}} f' \circ \Phi^{-1} + |y|^{-\frac{n+2}{2}} f'' \circ \Phi^{-1}.$$

Thus

$$\begin{aligned} \Delta_{\tilde{g}}(|y|^{-\frac{n+2}{2}} \tilde{f}) &= |y|^{-\frac{n+2}{2}} f'' \circ \Phi^{-1} - \left(\frac{n-2}{2}\right)^2 |y|^{-\frac{n+2}{2}} f \circ \Phi^{-1} + O(\varepsilon^2 |y|^{\frac{2-n}{2}}) f \circ \Phi^{-1} \\ &\quad + O(\varepsilon^2 |y|^{\frac{2-n}{2}}) f' \circ \Phi^{-1} \end{aligned}$$

which implies that

$$e^{-\frac{n+2}{2}t} \Delta_{\tilde{g}}(|y|^{\frac{2-n}{2}} \tilde{f}) \circ \Phi = f'' + O(\varepsilon^2 e^{-2t}) f' - \left(\left(\frac{n-2}{2}\right)^2 + O(\varepsilon^2 e^{-2t})\right) f.$$

Therefore,

$$\mathcal{L}_{\tilde{g}} f = f'' + O(\varepsilon^2 e^{-2t}) - f' \left(\left(\frac{n-2}{2}\right)^2 + O(\varepsilon^2 e^{-2t}) \right) f,$$

for any function f that depends only on t . Besides by (4.5) we obtain

$$B_{ij} = O(\varepsilon^2 e^{-2t}).$$

Thus, if $f = f_1 + f_2$, then

$$\mathbb{L}_{\tilde{g}}(f_1, f_2) = f'' + O(\varepsilon^2 e^{-2t}) f' - \left(\left(\frac{n-2}{2}\right)^2 + O(\varepsilon^2 e^{-2t})\right) f + O(\varepsilon^2 e^{-2t}) f. \quad (4.18)$$

Given a small number $\gamma > 0$, consider the linear operator \bar{L} given by

$$\bar{L}(f) = f'' + \gamma f' - \left(\left(\frac{n-2}{2}\right)^2 + \gamma\right) f. \quad (4.19)$$

and let $\gamma_1 = \frac{8-n}{2}\gamma > 0$ and $a(n) = \frac{1}{2(4-n)-\gamma_1}$.

We know proceed to the construction of the auxiliary functions. Define

$$h_\lambda(t) = \frac{a(n)}{2} c_1 \varepsilon^2 e^{(2-n)\lambda} e^{\frac{n-6}{2}t} (1 - e^{(4-n-\gamma_2)(t-\lambda)}), \quad (4.20)$$

where $\gamma_2 > 0$ is chosen such that the function $e^{(\frac{2-n}{2}-\gamma_2)t}$ is in the kernel of (4.19). Observe that we can make γ_1 and γ_2 as small as necessary depending on γ .

Note that the first two properties follows directly from the definition and moreover $h'_\lambda \leq 0$ in $(-\infty, \lambda]$. Since $t \geq -\log(\delta\varepsilon^{-1})$ it holds $\varepsilon^2 e^{-2t} \leq \delta^2$. Choosing $\delta^2 \ll \gamma$, if $f \geq 0$ and $f'_i \leq 0$, then by (4.18) and (4.19) we get

$$\mathbb{L}_{\hat{g}}(f_1, f_2) \geq f'' + \gamma f' - \left(\left(\frac{n-2}{2} \right)^2 + \gamma \right) f = \bar{L}(f).$$

Now observe that

$$h_\lambda(t) = a(n)c_1\varepsilon^2 e^{(2-n)\lambda} (e^{\frac{n-6}{2}t} - e^{(\frac{2-n}{2}-\gamma_2)t} e^{(4-n-\gamma_2)\lambda})$$

and

$$\begin{aligned} \bar{L}(h_\lambda) &= a(n)c_1\varepsilon^2 e^{(2-n)\lambda} \left(\left(\frac{n-6}{2} \right)^2 + \gamma \frac{n-6}{2} - \left(\frac{n-2}{2} \right)^2 - \gamma \right) e^{\frac{n-6}{2}t} \\ &= a(n)c_1\varepsilon^2 e^{(2-n)\lambda} \left(2(4-n) + \gamma \frac{n-8}{2} \right) e^{\frac{n-6}{2}t} = q_\lambda(t). \end{aligned}$$

Therefore

$$\mathbb{L}_{\hat{g}}(h_\lambda, h_\lambda) \geq \bar{L}(h_\lambda) = q_\lambda \geq |Q_\lambda|.$$

which proves (4.15). Besides,

$$h_\lambda(-\log(\delta\varepsilon^{-1})) = a(n)c_1 \left(\delta^{\frac{6-n}{2}} e^{(2-n)\lambda} \varepsilon^{\frac{n-2}{2}} - \delta^{\frac{n-2}{2}+\gamma_2} e^{(-2+\gamma_2)\lambda} \varepsilon^{\frac{6-n}{2}-\gamma_2} \right), \quad (4.21)$$

gives us the estimate (4.17).

Since $h'_\lambda \leq 0$, by (4.21) we obtain that

$$\max_{\Gamma_\lambda} h_\lambda \rightarrow 0 \text{ as } \lambda \rightarrow \infty. \quad (4.22)$$

For sufficiently large t_0 consider the set $\Gamma_{t_0} = [-\log(\delta\varepsilon^{-1}), t_0] \times \mathbb{S}^{n-1}$. Consequently we have that

$$v_{i,\lambda}(t) \leq C e^{\frac{2-n}{2}(2\lambda-t)},$$

forasmuch as $v_i = O(e^{\frac{n-2}{2}t})$. Define

$$w_{i,\lambda} = v_i - v_{i,\lambda} - h_\lambda \geq 0.$$

By (4.22) it follows that $w_{i,\lambda} \geq 0$ on Γ_{t_0} for sufficiently large λ . Let us show that $w_{i,\lambda} \geq 0$ for all $t \in [t_0, \lambda]$. A direct computation gives

$$h'_\lambda(t) = \frac{n-6}{4} a(n)c_1\varepsilon e^{(2-n)\lambda} e^{\frac{n-6}{2}t} \left(1 - \frac{2}{n-6} \left(\frac{2-n}{2} - \gamma_2 \right) e^{(4-n-\gamma_2)(t-\lambda)} \right).$$

Hence, for all $t \in [t_0, \lambda]$, we have

$$|h'_\lambda(t)| \leq C(\delta, \varepsilon, t_0)(e^{(2-n)\lambda} + e^{(-2+\gamma_2)\lambda}).$$

Now

$$\partial_t w_{i,\lambda}(t, \theta) = \partial_t v_i(t, \theta) + \partial_t v_i(2\lambda - t, \theta) - h'_\lambda(t). \quad (4.23)$$

Thus, because of the previous estimates and (4.9), when $t \in [t_0, \lambda]$, we get

$$\begin{aligned} \partial_t w_{i,\lambda}(t, \theta) &\leq -Ce^{\frac{2-n}{2}t} - Ce^{\frac{2-n}{2}(2\lambda-t)} - C(\delta, \varepsilon, t_0)(e^{(2-n)\lambda} + e^{(-2+\gamma_2)\lambda}) \\ &\leq -Ce^{\frac{2-n}{\lambda}} < 0, \end{aligned}$$

for t_0 and λ sufficiently large. Since $w_{i,\lambda}(\lambda, \theta) = 0$ for all $\theta \in \mathbb{S}^{n-1}$, we conclude that $w_{i,\lambda} \geq 0$ for all $t \in [t_0, \lambda]$, for sufficiently large λ . This finishes the proof of the Claim 2.

Once we proved the previous claim, using the same notation, we note that

$$w_{i,\lambda}(\lambda, \theta) = 0, \text{ for all } \theta \in \mathbb{S}^{n-1}$$

and

$$\mathbb{L}_{\hat{g}}(w_{1,\lambda}, w_{2,\lambda}) + \sum_{i=1}^2 b_{i,\lambda} w_{i,\lambda} = Q_\lambda - \mathbb{L}_{\hat{g}}(h_\lambda, h_\lambda) \leq 0.$$

In order to conclude the proof of Theorem 4.1, we will also need the following:

Claim 3: There exist $\lambda_0 > -\log(3n)$ and $\theta_0 \in \mathbb{S}^{n-1}$ such that

$$w_{\lambda_0}(-\log(\delta\varepsilon^{-1}), \theta_0) = 0.$$

Define

$$\lambda_0 = \inf\{\lambda_1; w_\lambda(t, \theta) \geq 0 \text{ in } \Gamma_\lambda, \forall \lambda \geq \lambda_1\}.$$

Note that (4.16) implies that this set is not empty. Besides, if we take $\lambda = -\log(3n)$, then by (4.9) and (4.23) we get

$$\partial_t w_\lambda(-\log(3n), \theta) = 2\partial_t v(-\log(3n), \theta) - 2h'_\lambda(-\log(3n)) > 0.$$

Moreover since $w_\lambda(-\log(3n), \theta) = 0$, we have $\lambda_0 > -\log(3n)$.

By continuity, $w_{\lambda_0} \geq 0$ in Γ_{λ_0} . Suppose the claim is false. Then $w_{\lambda_0}(-\log(\delta\varepsilon^{-1}), \theta) > 0$ for all $\theta \in \mathbb{S}^{n-1}$. We notice that by, (4.12), (4.5) and (4.15) we can apply the Maximum Principle, since for $\varepsilon > 0$ small enough we have

$$\Delta_{\hat{g}} w_{\lambda_0} - Dw_{\lambda_0} \leq \mathbb{L}_{\hat{g}}(w_{1,\lambda_0}, w_{2,\lambda_0}) = Q_{\lambda_0} - \mathbb{L}_{\hat{g}}(h_{\lambda_0}, h_{\lambda_0}) - \sum_{i=1}^2 b_{i,\lambda} w_{i,\lambda_0} \leq 0, \quad (4.24)$$

where D is a positive function. This implies that $w_{\lambda_0}(t, \theta) > 0$ for every $-\log(\delta\varepsilon^{-1}) < t < \lambda_0$ and $\theta \in \mathbb{S}^{n-1}$, since on the boundary $\partial\Gamma_{\lambda_0}$ we have $w_{\lambda_0} \geq 0$.

From definition of λ_0 , there exist a sequence $\{\lambda_j\}$ such that $\lambda_j < \lambda_0$ and $\lambda_j \rightarrow \lambda_0$, and a sequence $\{(t_j, \lambda_j)\}$ of interior minimum points of w_{λ_j} such that $(t_j, \theta_j) \rightarrow (t^*, \theta^*)$ with $w_{\lambda_j}(t_j, \theta_j) < 0$. Taking the limit we get $w_{\lambda_0}(t^*, \theta^*) = 0$ and $\nabla w_{\lambda_0}(t^*, \theta^*) = 0$. Therefore $t^* = \lambda_0$, but this is a contradiction to the Hopf's lemma. This proves the Claim 3.

With these claims on hand, let us prove the Theorem 4.1. By Claim 3, there exist $\lambda_0 > -\log(3n)$ and $\theta_0 \in \mathbb{S}^{n-1}$ such that

$$w_{\lambda_0}(-\log(\delta\varepsilon^{-1}), \theta_0) = 0.$$

Then, by definition of w_{λ_0} and (4.9) we get

$$0 < c(\delta) \leq v(-\log(\delta\varepsilon^{-1}), \theta_0) = (v_{\lambda_0} + 2h_{\lambda_0})(-\log(\delta\varepsilon^{-1}), \theta_0).$$

But $v(t, \theta) = O(e^{\frac{2-n}{2}(2\lambda-t)})$ implies that

$$v(-\log(\delta\varepsilon^{-1}), \theta_0) \leq c(\lambda_0, \delta)\varepsilon^{\frac{n-2}{2}}$$

and so

$$0 < c(\delta) \leq c(\lambda_0, \delta)\varepsilon^{\frac{n-2}{2}} + 2h_{\lambda_0}(-\log(\delta\varepsilon^{-1}), \theta_0). \quad (4.25)$$

But by (4.17) we obtain

$$h_{\lambda_0}(-\log(\delta\varepsilon^{-1}), \theta_0) \leq c\varepsilon^{\frac{1}{2}}$$

for $n = 3$,

$$h_{\lambda_0}(-\log(\delta\varepsilon^{-1}), \theta_0) \leq c\varepsilon^{1-\gamma}$$

for $n = 4$ and

$$h_{\lambda_0}(-\log(\delta\varepsilon^{-1}), \theta_0) \leq c\varepsilon^{\frac{1}{2}-\gamma}$$

for $n = 5$, which contradicts (4.25), since we can take the limit $\varepsilon \rightarrow 0$. This completes the proof of the theorem. ■

We may note that in the proof above we are not using that $-A$ is cooperative. As a consequence of the upper bound we get the following *spherical Harnack inequality*.

Corollary 4.2 Suppose \mathcal{U} is a positive smooth solution of (1) in $\Omega = B_1^n(0) \setminus \{0\}$, $3 \leq n \leq 5$, and assume that the potential A satisfies (H1). Then there exists a constant $c_1 > 0$ such that

$$\max_{|x|=r} u_i \leq c_1 \min_{|x|=r} u_i \quad (4.26)$$

for every $0 < r < \frac{1}{4}$. Moreover, $|\nabla u_i| \leq c_1 |x|^{-1} u_i$ and $|\nabla^2 u_i| \leq c_1 |x|^{-2} u_i$.

Proof. Define $u_{i,r}(y) = r^{\frac{n-2}{2}} u_i(ry)$, for every $0 < r < \frac{1}{4}$ and $|y| < r^{-1}$. Then the upper bound given by theorem (4.1) implies that $u_{i,r}(y) \leq c |y|^{\frac{2-n}{2}}$, for $|y| < \frac{1}{2} r^{-1}$. In particular, if $\frac{1}{2} \leq |y| \leq \frac{3}{2}$, we have that $u_{i,r}(y) \leq 2^{\frac{n-2}{2}} c$.

Moreover

$$\Delta_{g_r} u_{i,r}(y) - r^2 \sum_{j=1}^2 A_{ij}(ry) u_{j,r}(y) + \frac{n(n-2)}{4} |\mathcal{U}_r(y)|^{\frac{4}{n-2}} u_{i,r}(y) = 0.$$

where $(g_r)_{ij}(y) = g_{ij}(ry)$, which implies that

$$\Delta_{g_r} u_{i,r}(y) - r^2 A_{ii}(ry) u_{i,r}(y) = r^2 A_{ij}(ry) u_{j,r}(y) - \frac{n(n-2)}{4} |\mathcal{U}_r(y)|^{\frac{4}{n-2}} u_{i,r}(y).$$

for $j \neq i$. Using that $-A$ is cooperative, the Harnack inequality for linear elliptic equations and standard elliptic theory imply there exists $c_1 > 0$, not depending on r , such that

$$\max_{|x|=1} u_{i,r} \leq c_1 \min_{|x|=1} u_{i,r},$$

and $|\nabla u_{i,r}| + |\nabla^2 u_{i,r}| \leq c_1 u_{1,r}$ on the sphere of radius 1. This finishes the proof of the corollary. ■

4.2 Pohozaev invariants and removable singularities

In this section we will define the Pohozaev invariant of a solution and prove a removable singularity theorem for the solutions of the system (1). As a consequence we will derive a fundamental lower bound near the isolated singularity.

Given a positive solution \mathcal{U} to the system (1), we can define $P(r, \mathcal{U})$ similarly to (2.8) by

$$P(r, \mathcal{U}) = \int_{\partial B_r} \left(\frac{n-2}{2} \left\langle \mathcal{U}, \frac{\partial \mathcal{U}}{\partial \nu} \right\rangle - \frac{r}{2} |\nabla \mathcal{U}|^2 + r \left| \frac{\partial \mathcal{U}}{\partial \nu} \right|^2 + r \frac{(n-2)^2}{8} |\mathcal{U}|^{\frac{2n}{n-2}} \right) d\sigma.$$

The following lemma gives the Pohozaev-type identity we are interested in.

Lemma 4.3 (Pohozaev Identity) *Given $0 < s \leq r < 1$, it holds*

$$P(r, \mathcal{U}) - P(s, \mathcal{U}) = - \sum_{i=1}^2 \int_{B_r \setminus B_s} \left(x \cdot \nabla u_i + \frac{n-2}{2} u_i \right) \left((\Delta_g - \Delta) u_i - \sum_{j=1}^2 A_{ij}(x) u_j \right).$$

Proof. Adding and subtracting Δu_i in each system equation, we have

$$- \Delta u_i - \frac{n(n-2)}{4} |\mathcal{U}|^{\frac{4}{n-2}} u_i = (\Delta_g - \Delta) u_i - \sum_{j=1}^2 A_{ij}(x) u_j \quad (4.27)$$

in the punctured ball $\Omega = B_1^n(0) \setminus \{0\}$. Multiplying both sides by $x \cdot \nabla u_i$ and integrating over $B_r \setminus B_s$ we have

$$- \int_{B_r \setminus B_s} x \cdot \nabla u_i \left(\Delta u_i + \frac{n(n-2)}{4} |\mathcal{U}|^{\frac{4}{n-2}} u_i \right) = \int_{B_r \setminus B_s} x \cdot \nabla u_i \left((\Delta_g - \Delta)(u_i) - \sum_j A_{ij}(x) u_j \right).$$

First using integration by parts we get

$$\begin{aligned} \int_{B_r \setminus B_s} x \cdot \nabla u_i \Delta u_i &= \frac{n-2}{2} \int_{B_r \setminus B_s} |\nabla u_i|^2 - \frac{r}{2} \int_{\partial B_r} |\nabla u_i|^2 + \frac{s}{2} \int_{\partial B_s} |\nabla u_i|^2 \\ &+ r \int_{\partial B_r} \left| \frac{\partial u_i}{\partial \nu} \right|^2 - s \int_{\partial B_s} \left| \frac{\partial u_i}{\partial \nu} \right|^2. \end{aligned} \quad (4.28)$$

On the other hand, multiplying each equation for u_i and integrating over $B_r \setminus B_s$ we have

$$\begin{aligned} \int_{B_r \setminus B_s} u_i \left((\Delta_g - \Delta)(u_i) - \sum_{j=1}^2 A_{ij}(x) u_j \right) &= - \int_{B_r \setminus B_s} u_i \left(\Delta u_i + \frac{n(n-2)}{4} |\mathcal{U}|^{\frac{4}{n-2}} u_i \right) = \\ &= \int_{B_r \setminus B_s} |\nabla u_i|^2 - \int_{\partial B_r} u_i \frac{\partial u_i}{\partial \nu} + \int_{\partial B_s} u_i \frac{\partial u_i}{\partial \nu} - \frac{n(n-2)}{4} \int_{B_r \setminus B_s} |\mathcal{U}|^{\frac{4}{n-2}} u_i^2 \end{aligned}$$

which implies that

$$\begin{aligned} \int_{B_r \setminus B_s} |\nabla u_i|^2 &= \int_{\partial B_r} u_i \frac{\partial u_i}{\partial \nu} - \int_{\partial B_s} u_i \frac{\partial u_i}{\partial \nu} + \frac{n(n-2)}{4} \int_{B_r \setminus B_s} |\mathcal{U}|^{\frac{4}{n-2}} u_i^2 \\ &+ \int_{B_r \setminus B_s} u_i \left((\Delta_g - \Delta)(u_i) - \sum_{j=1}^2 A_{ij}(x) u_j \right). \end{aligned} \quad (4.29)$$

Replacing (4.29) in (4.28), we conclude that

$$\begin{aligned} &\int_{B_r \setminus B_s} \left(x \cdot \nabla u_i + \frac{n-2}{2} u_i \right) \left((\Delta_g - \Delta)(u_i) - \sum_{j=1}^2 A_{ij}(x) u_j \right) \\ &= - \int_{\partial B_r} \left[\frac{n-2}{2} u_i \frac{\partial u_i}{\partial \nu} - \frac{r}{2} |\nabla u_i|^2 + r \left| \frac{\partial u_i}{\partial \nu} \right|^2 \right] + \int_{\partial B_s} \left[\frac{n-2}{2} u_i \frac{\partial u_i}{\partial \nu} - \frac{s}{2} |\nabla u_i|^2 + s \left| \frac{\partial u_i}{\partial \nu} \right|^2 \right] \\ &\quad - \frac{n(n-2)^2}{8} \int_{B_r \setminus B_s} |\mathcal{U}|^{\frac{2n}{n-2}} - \frac{n(n-2)}{4} \int_{B_r \setminus B_s} (x \cdot \nabla u_i |\mathcal{U}|^{\frac{4}{n-2}} u_i). \end{aligned}$$

Note also that, using $\frac{n-2}{2n}\partial_k(|\mathcal{U}|^{\frac{2n}{n-2}}) = \sum_i |\mathcal{U}|^{\frac{4}{n-2}} u_i \partial_k u_i$, we obtain

$$\begin{aligned} & - \sum_{i=1}^2 \int_{B_r \setminus B_s} x \cdot \nabla u_i |\mathcal{U}|^{\frac{4}{n-2}} u_i = \frac{n-2}{2} \int_{B_r \setminus B_s} |\mathcal{U}|^{\frac{2n}{n-2}} - r \frac{n-2}{2n} \int_{\partial B_r} |\mathcal{U}|^{\frac{2n}{n-2}} \\ & + s \frac{n-2}{2n} \int_{\partial B_s} |\mathcal{U}|^{\frac{2n}{n-2}}, \end{aligned}$$

which finishes the proof. ■

In the case of the limit system we saw in 2.7 that $P(r, \mathcal{U})$ does not depend on r , and therefore is an invariant of the solution \mathcal{U} .

In order to define the invariant in a Riemannian setting, we need the upper bounds given by Theorem 4.1. In fact, since $g_{ij} = \delta_{ij} + O(|x|^2)$, we will have

$$\left| \left(x \cdot \nabla u_i + \frac{n-2}{2} u_i \right) \left((\Delta - \Delta_g) u_i - \sum_{j=1}^2 A_{ij}(x) u_j \right) \right| \leq c |x|^{2-n}, \quad (4.30)$$

for each coordinate i and the Pohozaev identity tell us the limit

$$P(\mathcal{U}) := \lim_{r \rightarrow 0} P(r, \mathcal{U})$$

exists. The number $P(\mathcal{U})$ is called the *Pohozaev invariant* of the solution $\mathcal{U} = (u_1, u_2)$.

Our main result of this subsection is the following.

Theorem 4.4 *Let \mathcal{U} be a positive solution to the system (1) in $B_1^n(0) \setminus \{0\}$, $3 \leq n \leq 5$ and that the potential A satisfies (H1) and (H2). Then $P(\mathcal{U}) \leq 0$. Moreover, $P(\mathcal{U}) = 0$ if and only if each coordinate u_i is smooth on the origin.*

The strategy of our proof will be assuming that $P(\mathcal{U}) \geq 0$ and then proving that in this case the origin is a removable singularity, and hence that the invariant is zero. In what follows let us denote by

$$u(x) = u_1(x) + u_2(x),$$

\bar{u} the average of u over ∂B_r , that is,

$$\bar{u}(r) := \oint_{\partial B_r} u := \frac{1}{\text{vol}(\partial B_r)} \int_{\partial B_r} u,$$

and define

$$w(t) = \bar{u}(r) r^{\frac{n-2}{2}},$$

where $t = -\ln r$.

We have divided the proof into a sequence of lemmas.

Lemma 4.5 Suppose \mathcal{U} be a positive solution of (1) which satisfies $P(\mathcal{U}) \geq 0$. Then

$$\liminf_{x \rightarrow 0} u(x)|x|^{\frac{n-2}{2}} = 0.$$

Proof. If this result is not true, without loss of generality we can suppose that there exist positive constants c_1, c_2 such that

$$c_1|x|^{\frac{2-n}{2}} \leq u_1(x) \leq c_2|x|^{\frac{2-n}{2}}, \quad (4.31)$$

where the second inequality above follows from Theorem 4.1. Choose any sequence $r_k \rightarrow 0$, and define

$$u_{i,k}(x) = r_k^{\frac{n-2}{2}} u_i(r_k x).$$

Then, using (4.31), we have the following estimate for $u_{1,k}$,

$$c_1|x|^{\frac{2-n}{2}} \leq u_{1,k}(x) \leq c_2|x|^{\frac{2-n}{2}}. \quad (4.32)$$

Moreover $u_{i,k}$ satisfies

$$-\Delta_{g_k} u_{i,k} + r_k^2 \sum_{j=1}^2 A_{ij}(r_k x) u_{j,k} = \frac{n(n-2)}{4} |\mathcal{U}_k|^{\frac{4}{n-2}} u_{i,k} \quad \text{in } B_{r_k^{-1}}(0) \setminus \{0\},$$

where $(g_k)_{lm}(x) = g_{lm}(r_k x)$. Elliptic theory then implies that there exists a subsequence, also denoted by $u_{i,k}$, which converges in compact subsets of $\mathbb{R}^n \setminus \{0\}$ to a solution $\mathcal{U}_0 = (u_{1,0}, u_{2,0})$ of the limit system

$$\Delta u_{i,0} + \frac{n(n-2)}{4} |\mathcal{U}_0|^{\frac{4}{n-2}} u_{i,0} = 0.$$

By (4.32) we get

$$u_{1,0}(x) \geq c_1|x|^{\frac{2-n}{2}},$$

which implies that \mathcal{U}_0 is a singular at the origin. However, by Theorem 2.8 we know that $P(\mathcal{U}_0) < 0$. This is a contradiction, because

$$P(\mathcal{U}_0) = P(1, \mathcal{U}_0) = \lim_{k \rightarrow \infty} P(1, \mathcal{U}_k) = \lim_{k \rightarrow \infty} P(r_k, \mathcal{U}) = P(\mathcal{U}) \geq 0.$$

■

Lemma 4.6 Assume that \mathcal{U} is a positive solution of (1) and that the potential A satisfies (H1) and (H2). Suppose that there exists a sequence $(t_k)_{k \in \mathbb{N}}$ of minimum points for

w such that $\lim_{k \rightarrow \infty} w(t_k) = 0$. Along $|x| = r_k$, where $t_k = -\ln r_k$, there exist positive constants a_i, b_i such that

$$\begin{aligned} u_i(x) &= \bar{u}(r_k)(c_i + o(1)) \\ |\nabla u_i(x)| &= -\bar{u}'(r_k)(2a_i + o(1)). \end{aligned} \tag{4.33}$$

where $c_i = a_i + b_i$. Moreover $\langle a, b \rangle \neq 0$.

Proof. Let $r_k = e^{-t_k}$ and define $w_i = r^{\frac{n-2}{2}} \bar{u}_i$ and

$$v_{i,k}(y) = r_k^{\frac{n-2}{2}} u_i(r_k y).$$

Once that $\bar{v}_{i,k}(1) = w_1(t_k) \leq w_1(t_k) + w_2(t_k) = w(t_k) \rightarrow 0$ we get from Harnack inequality that each coordinate $v_{i,k}$ converge uniformly in compact subsets of $\mathbb{R}^n \setminus \{0\}$ to 0. So, if we define

$$h_{i,k}(y) = v_k(p)^{-1} v_{i,k}(y)$$

where $v_k = \sum_i v_{i,k}$ and $p = (1, 0, \dots, 0)$, we have that

$$\begin{aligned} -\Delta_{g_k} h_{i,k}(y) &= -v_k(p)^{-1} (\Delta_{g_k} v_{i,k}) \\ &= v_k(p)^{-1} \left(\frac{n(n-2)}{4} |\mathcal{V}_k|^{\frac{4}{n-2}} v_{i,k} - r_k^2 \sum_{j=1}^2 A_{ij}(r_k y) v_{j,k} \right). \end{aligned}$$

This implies

$$-\Delta_{g_k} h_{i,k} + r_k^2 \sum_{j=1}^2 A_{ij}(r_k y) h_{j,k} = \frac{n(n-2)}{4} v_k(p)^{\frac{4}{n-2}} |H_k|^{\frac{4}{n-2}} h_{i,k},$$

where $(g_i)_{lm}(y) = g_{lm}(r_i y)$ and $H_k = (h_{1,k}, h_{2,k})$. By elliptic estimates we know that there exists a subsequence $h_{i,k}$ which converge in C_{loc}^2 to a nonnegative harmonic function h_i in $\mathbb{R}^n \setminus \{0\}$. Then

$$h_i(y) = a_i |y|^{2-n} + b_i,$$

and $a_1 + a_2 = b_1 + b_2 = \frac{1}{2}$ since $h_1(p) + h_2(p) = 1$ and $\partial_r((h_1 + h_2)(r) r^{\frac{n-2}{2}}) = 0$ at $r = 1$.

Now, multiplying line i of system (1) by u_j and integrating over $B_{r_k}(0) \setminus B_{\varepsilon r_k}(0)$, we obtain that

$$\int_{B_{r_k} \setminus B_{\varepsilon r_k}} u_j \Delta_g u_i dv_g + \sum_{l=1}^2 \int_{B_{r_k} \setminus B_{\varepsilon r_k}} A_{il} u_l dv_g = \frac{n(n-2)}{4} \int_{B_{r_k} \setminus B_{\varepsilon r_k}} |\mathcal{U}|^{\frac{4}{n-2}} u_i u_j dv_g.$$

Reversing i and j and subtracting one equation from the other, we obtain

$$\int_{B_{r_k} \setminus B_{\varepsilon r_k}} (u_j \Delta_g u_i - u_i \Delta_g u_j) dv_g = \sum_{l=1}^2 \int_{B_{r_k} \setminus B_{\varepsilon r_k}} (u_i A_{jl} - u_j A_{il}) u_l dv_g,$$

which, integrating by parts, guarantees that

$$\int_{\partial B_{r_k}} (u_j \partial_r u_i - u_i \partial_r u_j) d\sigma_g = \sum_{l=1}^2 \int_{B_{r_k} \setminus B_{\varepsilon r_k}} (u_i A_{jl} - u_j A_{il}) u_l dv_g + \int_{\partial B_{\varepsilon r_k}} (u_j \partial_r u_i - u_i \partial_r u_j) d\sigma_g. \quad (4.34)$$

In order to analyse the last integral on the right-hand side, let

$$\varphi_{i,k}^\varepsilon(z) = (\varepsilon r_k)^{\frac{n-2}{2}} u_i(\varepsilon r_k z).$$

Then

$$\int_{\partial B_{\varepsilon r_k}} (u_j \partial_r u_i - u_i \partial_r u_j) d\sigma_g = \int_{\partial B_1} (\varphi_{j,k}^\varepsilon \partial_r \varphi_{i,k}^\varepsilon - \varphi_{i,k}^\varepsilon \partial_r \varphi_{j,k}^\varepsilon) d\sigma_g.$$

On the other hand

$$-\Delta_{g_{\varepsilon k}} \varphi_{i,k}^\varepsilon + (\varepsilon r_k)^2 \sum_{j=1}^2 \tilde{A}_{ij}^\varepsilon \varphi_{j,k}^\varepsilon = c(n) |\varphi_k^\varepsilon|^{\frac{4}{n-2}} \varphi_{i,k}^\varepsilon$$

in $B_{(\varepsilon r_k)^{-1}}(0) \setminus \{0\}$ and by theorem (4.1)

$$|\varphi_k^\varepsilon(z)| \leq C |z|^{\frac{2-n}{2}}.$$

Similarly to what we did at the beginning of this proof, after passing to a subsequence, $\varphi_{i,k}^\varepsilon$ converges in C^2 topology locally in compact subsets of $\mathbb{R}^n \setminus \{0\}$, to a positive solution

$$\Delta u_i + \frac{n(n-2)}{4} |\mathcal{U}|^{\frac{4}{n-2}} u_i = 0$$

which using Theorem (2.10) as well Proposition 1.1 in [7] is of the form $u_0 \Lambda$, where $\Lambda = (\Lambda_1, \Lambda_2)$ is a vector in the unit sphere with nonnegative coordinates. Consequently, when ε goes to zero we have

$$\lim_{\varepsilon \rightarrow 0} \int_{\partial B_1} (\varphi_{j,k}^\varepsilon \partial_r \varphi_{i,k}^\varepsilon - \varphi_{i,k}^\varepsilon \partial_r \varphi_{j,k}^\varepsilon) d\sigma_g = \int_{\partial B_1} (u_0 \partial_r u_0 \Lambda_i \Lambda_j - u_0 \partial_r u_0 \Lambda_j \Lambda_i) d\sigma_g = 0.$$

Making $\varepsilon \rightarrow 0$ in (4.34), we conclude that

$$\int_{\partial B_{r_k}} (u_j \partial_r u_i - u_i \partial_r u_j) d\sigma_g = \sum_{l=1}^2 \int_{B_{r_k}} (u_i A_{jl} - u_j A_{il}) u_l dv_g. \quad (4.35)$$

Using the same notation from the first claim, by the hypothesis under the potential A (H2) and by Theorem 4.1 the equation above implies that

$$\begin{aligned} & \int_{\partial B_1} (h_{j,k} \partial_r h_{i,k} - h_{i,k} \partial_r h_{j,k}) d\sigma_g \\ &= r_k^2 \sum_{l=1}^2 \int_{B_1} \frac{(u_i(r_k x) A_{jl}(r_k x) - u_j(r_k x) A_{il}(r_k x)) u_l(r_k x)}{u(r_k p)^2} dv_g \\ &= O(r_k^{5-n} u(r_k p)^{-2}), \end{aligned}$$

which converges to zero when dimension is less or equal than five. Finally, we conclude that

$$\int_{\partial B_1} (h_j \partial_r h_i - h_i \partial_r h_j) d\sigma_g = 0,$$

and consequently

$$a_1 b_2 = a_2 b_1$$

which finishes the proof. \blacksquare

Lemma 4.7 *Assume that \mathcal{U} is a positive solution of (1) and along $|x| = r_k$ the solution satisfies (4.33). Then*

$$P(r_k, \mathcal{U}) = \sigma_{n-1} \left(-4\langle a, b \rangle \frac{\delta^2}{2} w^2(t_k) + \frac{\delta^2}{2} (c_1^2 + c_2^2)^{\frac{n}{n-2}} |W|^{\frac{2n}{n-2}}(t_k) \right) (1 + o(1)). \quad (4.36)$$

Proof. Since the solution \mathcal{U} satisfies (4.33), note that

$$\left\langle \mathcal{U}, \frac{\partial \mathcal{U}}{\partial \nu} \right\rangle = \sum_{i=1}^2 \bar{u}(r_k) \bar{u}'(r_k) (2c_i a_i + o(1)).$$

On the other hand, using that $w_t(t_k) = 0$ it holds the following equality

$$\bar{u}'(r_k) r_k^{\frac{n}{2}} = -\frac{n-2}{2} w(t_k). \quad (4.37)$$

Multiplying (4.37) by $\bar{u}(r_k)$, we conclude that

$$\frac{n-2}{2} \left\langle \mathcal{U}, \frac{\partial \mathcal{U}}{\partial \nu} \right\rangle = - \sum_{i=1}^2 r_k^{1-n} \frac{(n-2)^2}{2} w^2(t_k) (c_i a_i + o(1)).$$

Similarly, we also have

$$\begin{aligned} -\frac{r}{2} |\nabla \mathcal{U}|^2 + r \left| \frac{\partial \mathcal{U}}{\partial \nu} \right|^2 &= r_k \sum_{i=1}^2 \bar{u}'(r_k)^2 (2a_i^2 + o(1)) \\ &= r_k^{1-n} \sum_i \frac{(n-2)^2}{2} w^2(t_k) (a_i^2 + o(1)). \end{aligned}$$

With these two equalities in hands we get

$$\begin{aligned} P(r_k, \mathcal{U}) &= \int_{\partial B_{r_k}} \left(\frac{n-2}{2} \left\langle \mathcal{U}, \frac{\partial \mathcal{U}}{\partial \nu} \right\rangle - \frac{r}{2} |\nabla \mathcal{U}|^2 + r \left| \frac{\partial \mathcal{U}}{\partial \nu} \right|^2 + r \frac{(n-2)^2}{8} |\mathcal{U}|^{\frac{2n}{n-2}} \right) d\sigma \\ &= \sigma_{n-1} \left(-4\langle a, b \rangle \frac{\delta^2}{2} w(t_k)^2 + \frac{\delta^2}{2} (c_1^2 + c_2^2)^{\frac{n}{n-2}} |W|^{\frac{2n}{n-2}}(t_k) \right) (1 + o(1)), \end{aligned}$$

which finishes the proof. \blacksquare

Lemma 4.8 *Let \mathcal{U} a positive solution of (1) defined in the punctured ball. If*

$$\lim_{|x| \rightarrow 0} |x|^{\frac{n-2}{2}} u(x) = 0, \quad (4.38)$$

then \mathcal{U} extends as a smooth solution to all of B^n .

Proof. We begin by obtaining upper and lower bounds for the second derivatives of w in terms of w . Indeed, observe that the upper bound in Theorem 4.1 implies that $w(t)$ is bounded. Then,

$$\bar{u}_r = \oint_{\partial B_r} u_r,$$

and since

$$w_t = -\bar{u}_r r^{\frac{n}{2}} - \frac{n-2}{2} w,$$

we also get that $|w_t|$ is bounded. Derivating again the function w we obtain

$$w_{tt} = \bar{u}_{rr} r^{\frac{n+2}{2}} + (n-1) \bar{u}_r r^{\frac{n}{2}} + \left(\frac{n-2}{2} \right)^2 w. \quad (4.39)$$

Choosing a fixed $s < r$, by the Divergence Theorem, we get

$$\begin{aligned} \left(\int_{B_r \setminus B_s} \Delta u \right)_r &= \left(\int_{\partial B_r} u_r \right)_r = \left(\int_{\partial B_1} u_r(r \cdot) r^{n-1} \right)_r \\ &= \frac{n-1}{r} \int_{\partial B_1} u_r(r \cdot) r^{n-1} + \int_{\partial B_1} u_{rr}(r \cdot) r^{n-1} = \frac{n-1}{r} \int_{\partial B_r} u_r + r^{n-1} \sigma_{n-1} \left(\oint_{\partial B_r} u_r \right)_r, \end{aligned}$$

where $\sigma_{n-1} = \text{vol}(\mathbb{S}^{n-1})$. Thus

$$\begin{aligned} \bar{u}_{rr} &= \left(\oint_{\partial B_r} u_r \right)_r = (1-n)r^{-1} \oint_{\partial B_r} u_r + \sigma_{n-1}^{-1} r^{1-n} \left(\int_{B_r \setminus B_s} \Delta u \right)_r \\ &= \frac{1-n}{r} \bar{u}_r + \sigma_{n-1}^{-1} r^{1-n} \sum_{i=1}^2 \left(\int_{B_r \setminus B_s} \left((\Delta - \Delta_g) u_i + \sum_{j=1}^2 A_{ij} u_j - c(n) |\mathcal{U}|^{\frac{4}{n-2}} u_i \right) \right)_r \end{aligned}$$

and so

$$\bar{u}_{rr} + \frac{n-1}{r} \bar{u}_r = \oint_{\partial B_r} \left((\Delta - \Delta_g) u + \sum_{i,j=1}^2 A_{ij}(x) u_j - c(n) |\mathcal{U}|^{\frac{4}{n-2}} u \right)$$

where $c(n) = \frac{n(n-2)}{4}$. Replacing in (4.39) we have that

$$w_{tt} - \left(\frac{n-2}{2} \right)^2 w = r^{\frac{n+2}{2}} \oint_{\partial B_r} \left((\Delta - \Delta_g) u + \sum_{i,j=1}^2 A_{ij}(x) u_j - c(n) |\mathcal{U}|^{\frac{4}{n-2}} u \right).$$

Applying the spherical Harnack inequality in Corollary 4.2 for each coordinate function we have that

$$C^{-1}\bar{u}^{\frac{n+2}{n-2}} \leq c(n) \oint_{\partial B_r} |\mathcal{U}|^{\frac{4}{n-2}} u \leq C\bar{u}^{\frac{n+2}{n-2}}$$

and

$$\left| \oint_{\partial B_r} \left((\Delta - \Delta_g)u + \sum_{i,j=1}^2 A_{ij}(x)u_j \right) \right| \leq c\bar{u}.$$

With these estimates we obtain the following inequality

$$-c_1 w^{\frac{n+2}{n-2}} - c_3 e^{-2t} w \leq w_{tt} - \left(\frac{n-2}{2} \right)^2 w \leq -c_2 w^{\frac{n+2}{n-2}} + c_3 e^{-2t} w. \quad (4.40)$$

By hypothesis (4.38) we know that $\lim_{t \rightarrow \infty} w(t) = 0$. The strategy is to show that $u \in L_{loc}^p(B_1^n(0))$ for some $p > 2n/(n-2)$ and then by elliptic theory the function u extends smoothly across the origin. Consequently, each coordinate function u_i is smooth.

Note that the first inequality in (4.40) implies that there exists $\varepsilon_0 > 0$, such that if $w(t) \leq \varepsilon_0$ and t is sufficiently large, then $w_{tt}(t) > 0$. Since $\lim_{t \rightarrow \infty} w(t) = 0$, there exists T_1 so that $w(t) < \varepsilon_0$ and $w_{tt} > 0$ for $t \geq T_1$. This implies that $w_t < 0$ for $t \geq T_1$.

By the first inequality in (4.40), given any positive number $0 < \delta < n-2$, there exists T_0 sufficiently large such that

$$w_{tt} - \left(\frac{n-2}{2} - \delta \right)^2 w \geq (\delta(n-2) - \delta^2 - c_1 w^{\frac{4}{n-2}} - c_3 e^{-2t})w \geq 0$$

for $t \geq T_0$, which implies that

$$\left(w_t^2 - \left(\frac{n-2}{2} - \delta \right)^2 w^2 \right)_t = 2w_t \left(w_{tt} - \left(\frac{n-2}{2} - \delta \right)^2 w \right) \leq 0$$

for $t > T_2 = \max\{T_0, T_1\}$, and using that $\lim_{t \rightarrow \infty} w_t(t) = 0$, we obtain

$$w_t^2 - \left(\frac{n-2}{2} - \delta \right)^2 w^2 \geq 0.$$

By integrating we get, for $t \geq T_2$, that

$$w(t) \leq w(T_0) e^{-\left(\frac{n-2}{2} - \delta\right)(t-T_0)}.$$

Equivalently, there exists $r_0(\delta) > 0$, so that

$$u(x) \leq c(\delta)|x|^{-\delta} \text{ for all } x \in B_{r_0}(0).$$

Since $\delta > 0$ is arbitrarily small, the estimate above implies that $u \in L_{loc}^p(B_1(0))$ for arbitrarily large p , which finishes our proof. ■

Now we are ready to prove the main result of this section.

Proof of Theorem 4.4. Following the aforementioned strategy let us suppose that $P(\mathcal{U}) \geq 0$. By Lemma 4.8 the proof is completed by showing that $\lim_{x \rightarrow 0} u(x)|x|^{\frac{n-2}{2}} = 0$. Suppose by contradiction that this is false.

Since the Pohozaev invariant is nonnegative, it holds as a consequence of Lemma 4.5 that $\liminf_{x \rightarrow 0} u(x)|x|^{\frac{n-2}{2}} = 0$. Then we will assume that $\limsup_{x \rightarrow 0} u(x)|x|^{\frac{n-2}{2}} > 0$. Hence we can choose $\varepsilon_0 > 0$ sufficiently small so that we are able to construct sequences $\bar{t}_k \leq t_k \leq t_k^*$ satisfying $\lim_{k \rightarrow \infty} \bar{t}_k = +\infty$, $w(\bar{t}_k) = w(t_k^*) = \varepsilon_0$, $w_t(t_k) = 0$ and $\lim_{k \rightarrow \infty} w(t_k) = 0$.

Using the results obtained on Lemmas 4.6 and 4.7, we have that

$$P(r_k, \mathcal{U}) = \sigma_{n-1} \left(-4\langle a, b \rangle \frac{\delta^2}{2} w^2(t_k) + \frac{\delta^2}{2} (c_1^2 + c_2^2)^{\frac{n}{n-2}} |W|^{\frac{2n}{n-2}}(t_k) \right) (1 + o(1)), \quad (4.41)$$

and consequently

$$P(\mathcal{U}) = \lim_{k \rightarrow \infty} P(r_k, \mathcal{U}) = 0.$$

Moreover, using that Lemma 4.6 guarantees that $\langle a, b \rangle \neq 0$ with the Pohozaev identity, we have the following estimate

$$w^2(t_k) \leq c|P(r_k, \mathcal{U})| \leq c(I_1 + I_2), \quad (4.42)$$

where

$$I_1 = \int_{B_{r_k} \setminus B_{r_k}^*} |A(\mathcal{U})| dx,$$

$$I_2 = \int_{B_{r_k}^*} |A(\mathcal{U})| dx,$$

and $A(\mathcal{U}) = \sum_i (x \cdot \nabla u_i + \frac{n-2}{2} u_i) \left((\Delta_g - \Delta)(u_i) - \sum_j A_{ij}(x) u_j \right)$. We will follow a series of calculations with the goal of obtaining better estimates for the term I_1 . First remember the inequality (4.40) obtained in the proof of Lemma 4.8

$$-c_1 w^{\frac{n+2}{n-2}} - c_3 e^{-2t} w \leq w_{tt} - \left(\frac{n-2}{2} \right)^2 w \leq -c_2 w^{\frac{n+2}{n-2}} + c_3 e^{-2t} w. \quad (4.43)$$

It follows by the first inequality of (4.43) that

$$w_{tt} - \left(\frac{n-2}{2} \right)^2 w \geq -c_1 w^{\frac{n+2}{n-2}} - c_3 e^{-2t} w \geq -c_1 w^{\frac{n+2}{n-2}} - c_3 e^{-2t_k} w$$

for $t \geq t_k$, which implies

$$\frac{d}{dt} \left[w_t^2 - \left(\left(\frac{n-2}{2} \right)^2 - c_3 e^{-2t_k} \right) w^2 + \frac{n-2}{n} c_1 w^{\frac{2n}{n-2}} \right] \geq 0$$

for $t_k \leq t \leq t_k^*$.

Hence, if $t_k \leq t \leq t_k^*$, then

$$w_t(t)^2 - g(w(t)) + g(w(t_k)) = \int_{t_k}^t \frac{d}{dt} (w_t^2 - g(w)) \geq 0,$$

which implies that

$$\frac{dw}{dt} \geq \sqrt{g(w) - g(w(t_k))}$$

and so

$$t - t_k = \int_{w(t_k)}^{w(t)} \frac{dt}{dw} dw \leq \int_{w(t_k)}^{w(t)} \frac{dw}{\sqrt{g(w) - g(w(t_k))}},$$

where

$$g(w) = \left(\left(\frac{n-2}{2} \right)^2 - c_3 e^{-2t_k} \right) w^2 - \frac{n-2}{n} c_1 w^{\frac{2n}{n-2}}.$$

Introducing the variable $\eta = \frac{w(t)}{w(t_k)}$, we get

$$t - t_k \leq \int_1^{\frac{w(t)}{w(t_k)}} \frac{d\eta}{\sqrt{\bar{g}(\eta) - \bar{g}(1)}} = \int_1^{\frac{w(t)}{w(t_k)}} \sqrt{\frac{\eta^2 - 1}{\bar{g}(\eta) - \bar{g}(1)}} \frac{d\eta}{\sqrt{\eta^2 - 1}}, \quad (4.44)$$

where

$$\bar{g}(\eta) = \left(\left(\frac{n-2}{2} \right)^2 - c_3 e^{-2t_k} \right) \eta^2 - \frac{n-2}{n} c_1 w(t_k)^{\frac{4}{n-2}} \eta^{\frac{2n}{n-2}}.$$

First, since $1 \leq \eta \leq \frac{w(t)}{w(t_k)} \leq \frac{\varepsilon_0}{w(t_k)}$, we have that

$$\frac{w(t_k)^{\frac{4}{n-2}} \left(\eta^{\frac{2n}{n-2}} - 1 \right)}{\eta^2 - 1} \leq c w(t_k)^{\frac{4}{n-2}} \eta^{\frac{4}{n-2}} \leq c \varepsilon_0^{\frac{4}{n-2}},$$

and we observe that

$$w(t_k)^{\frac{4}{n-2}} \int_1^{\frac{w(t)}{w(t_k)}} \frac{\eta^{\frac{4}{n-2}}}{\sqrt{\eta^2 - 1}} d\eta \leq c.$$

Now

$$\left(\frac{\eta^2 - 1}{\bar{g}(\eta) - \bar{g}(1)} \right)^{\frac{1}{2}} \leq \frac{2}{n-2} + c e^{-2t_k} + c \frac{w(t_k)^{\frac{4}{n-2}} \left(\eta^{\frac{2n}{n-2}} - 1 \right)}{\eta^2 - 1}.$$

Finally, since

$$\int_1^{\frac{w(t)}{w(t_k)}} \frac{d\eta}{\sqrt{\eta^2 - 1}} \leq c + \ln \frac{w(t)}{w(t_k)},$$

we obtain

$$\int_1^{\frac{w(t)}{w(t_k)}} \frac{d\eta}{\sqrt{\bar{g}(\eta) - \bar{g}(1)}} \leq \left(\frac{2}{n-2} + ce^{-2t_k} \right) \ln \frac{w(t)}{w(t_k)} + c.$$

From inequality (4.44), we get

$$t - t_k \leq \left(\frac{2}{n-2} + ce^{-2t_k} \right) \ln \frac{w(t)}{w(t_k)} + c \quad (4.45)$$

for all $t \in (t_k, t_k^*)$.

In order to estimate $t - t_k$ from below, we first observe that the second inequality in (4.43) implies that

$$w_{tt} - \left(\left(\frac{n-2}{2} \right)^2 + ce^{-2t_k} \right) w \leq 0.$$

Then the function $w_t^2 - \left(\left(\frac{n-2}{2} \right)^2 + ce^{-2t_k} \right) w^2$ is decreasing in (t_k, t_k^*) , and therefore

$$w_t^2 - \left(\left(\frac{n-2}{2} \right)^2 + ce^{-2t_k} \right) w^2 \leq - \left(\left(\frac{n-2}{2} \right)^2 + ce^{-2t_k} \right) w^2(t_k).$$

Hence

$$w_t \leq \sqrt{\left(\left(\frac{n-2}{2} \right)^2 + ce^{-2t_k} \right) (w^2 - w^2(t_k))},$$

and then

$$t - t_k = \int_{w(t_k)}^{w(t)} \frac{dt}{dw} dw \geq \left(\frac{2}{n-2} - ce^{-2t_k} \right) \int_{w(t_k)}^{w(t)} \frac{dw}{\sqrt{w^2 - w^2(t_k)}}.$$

Together with inequality (4.45), we get for $t_k \leq t \leq t_k^*$, that

$$\left(\frac{2}{n-2} - ce^{-2t_k} \right) \ln \frac{w(t)}{w(t_k)} \leq t - t_k \leq \left(\frac{2}{n-2} + ce^{-2t_k} \right) \ln \frac{w(t)}{w(t_k)} + c. \quad (4.46)$$

Similarly one can prove that, for $\bar{t}_k \leq t \leq t_k$, it holds

$$\left(\frac{2}{n-2} - ce^{-2\bar{t}_k} \right) \ln \frac{w(t)}{w(t_k)} \leq t_k - t \leq \left(\frac{2}{n-2} + ce^{-2\bar{t}_k} \right) \ln \frac{w(t)}{w(t_k)} + c. \quad (4.47)$$

Once we get the above inequalities, let's go back to the estimates of the terms I_1 and I_2 in inequality (4.42). Recall that by (4.30), $|A(\mathcal{U})| \leq c|x|^{2-n}$, and therefore

$$I_2 \leq c(r_k^*)^2 = ce^{-2t_k^*}.$$

From the first inequality in (4.46), we obtain

$$w(t) \leq w(t_k) \exp \left(\left(\frac{n-2}{2} + ce^{-2t_k} \right) (t - t_k) \right),$$

which implies

$$v(x) \leq cw(t_k) \exp \left(- \left(\frac{n-2}{2} + ce^{-2t_k} \right) t_k \right) r^{2-n-ce^{-2t_k}}. \quad (4.48)$$

Recall that, by the spherical Harnack inequality 4.2, for each coordinate function, $u_i \leq Cr^{\frac{2-n}{2}}$, $|\nabla u_i| \leq Cr^{-1}u_i$, and $|\nabla^2 u_i| \leq Cr^{-2}u_i$, so

$$|A(\mathcal{U})| \leq Cr^{\frac{2-n}{2}}u.$$

Using the estimate (4.48) we obtain

$$I_1 \leq cw(t_k)e^{-\frac{n-2}{2}t_k-ce^{-2t_k}t_k} \int_{B_{r_k} \setminus B_{r_k^*}} |x|^{3-\frac{3n}{2}-ce^{-2t_k}} dx,$$

and so

$$I_1 \leq cw(t_k)e^{-2t_k}.$$

Therefore, from (4.42) and the estimates for I_1 and I_2 , we get

$$w^2(t_k) \leq cw(t_k)e^{-2t_k} + ce^{-2t_k^*}.$$

Passing to subsequences, if necessary, we can suppose either

$$w^2(t_k) \leq cw(t_k)e^{-2t_k} \quad (4.49)$$

or

$$w^2(t_k) \leq ce^{-2t_k^*}. \quad (4.50)$$

Define $L_k = -\frac{2}{n-2} \log w(t_k)$ and choose $\delta > 0$ small. Then, from the first inequality in (4.47), we get

$$t_k - \bar{t}_k \geq (1 - \delta)L_k - c, \quad (4.51)$$

and adding to the first inequality in (4.46), we obtain

$$t_k^* - \bar{t}_k \geq (2 - 2\delta)L_k - c. \quad (4.52)$$

If inequality (4.49) holds, then $w(t_k) \leq ce^{-2t_k}$ and so $L_k \geq \frac{4}{n-2}t_k - c$. From inequality (4.51), we conclude

$$t_k - t_k^* \geq (1 - \delta)\frac{4}{n-2}t_k - c,$$

and consequently

$$\bar{t}_k \leq \left(\frac{n-6}{n-2} + \frac{4\delta}{n-2} \right) t_k + c.$$

The inequality above gives us a contradiction since $t_k^* \geq t_k \geq \bar{t}_k \rightarrow \infty$ as $k \rightarrow \infty$ and on the other hand $\frac{n-6}{n-2} + \frac{4\delta}{n-2} < 0$ by our assumption $3 \leq n \leq 5$.

If inequality (4.50) holds, then $L_k \geq \frac{2}{n-2} t_k^* + c$. From inequality (4.52), we get

$$\bar{t}_k \leq t_k^* - (2 - 2\delta)L_k + c,$$

and so

$$\bar{t}_k \leq \left(\frac{n-6}{n-2} + 2\delta \right) t_k^* + c.$$

If $3 \leq n \leq 5$, this is again a contradiction by the same reasons as before. Then we conclude that $\lim_{x \rightarrow 0} u(x)|x|^{\frac{n-2}{2}} = 0$, and the result follows as consequence of Lemma 4.8.

■

As a consequence of the removable singularity theorem, we can now establish a fundamental lower bound.

Corollary 4.9 *Assume $3 \leq n \leq 5$ and let \mathcal{U} a positive solution to the system (1) in $B_1^n(0) \setminus \{0\}$ and that the potential A satisfies (H1) and (H2). If 0 is a nonremovable singularity, then there exists $c > 0$ such that*

$$|\mathcal{U}|(x) \geq c d_g(x, 0)^{\frac{2-n}{2}}$$

for $0 < d_g(x, 0) < \frac{1}{2}$.

Proof. Suppose by contradiction that this is not true. Then $\liminf_{t \rightarrow \infty} w(t) = 0$, where $w(t) = r^{\frac{n-2}{2}} \bar{u}(r)$, $u = \sum_{i=1}^2 u_i$ and $t = -\log r$, as in proof of the Theorem 4.4. Since 0 is a nonremovable singularity, we also have $\limsup_{t \rightarrow \infty} w(t) > 0$, otherwise we contradict Lemma 4.8. Therefore there exists a sequence $t_k \rightarrow \infty$ such that $w_t(t_k) = 0$ and $\lim_{k \rightarrow \infty} w(t_k) = 0$. So, if $r_k = e^{-t_k}$ we can check by Lemmas 4.6 and 4.7

$$P(r_k, \mathcal{U}) = \sigma_{n-1} \left(-4 \langle a, b \rangle \frac{\delta^2}{2} w^2(t_k) + \frac{\delta^2}{2} (c_1^2 + c_2^2)^{\frac{n}{n-2}} |W|^{\frac{2n}{n-2}}(t_k) \right) (1 + o(1)).$$

where $|W|^2 = w_1^2 + w_2^2$. But, in this case

$$P(\mathcal{U}) = \lim_{i \rightarrow \infty} P(r_k, \mathcal{U}) = 0,$$

which is a contradiction. This finishes the proof. ■

4.3 Convergence to a Radial Solution

Our main goal in this section is to prove that a local singular solution to our system is asymptotic to a radial Fowler-type solution, near the nonremovable isolated singularity. The proof will follow exactly the same steps of the Theorem 3.7, however since we are dealing with an arbitrary Riemannian metric we need to take exponential correction terms into account in order to obtain analogous convergence results for the Jacobi fields.

Theorem 4.10 *Suppose that \mathcal{U} is a solution of the system (1) in the punctured ball $B_1^n(0) \setminus \{0\}$ and that the potential A satisfies (H1) and (H2). If there exist positive constants c_1 and c_2 such that*

$$c_1|x|^{\frac{2-n}{2}} \leq |\mathcal{U}|(x) \leq c_2|x|^{\frac{2-n}{2}} \quad (4.53)$$

then there exist a Fowler-type solution $\mathcal{U}_0 = u_0\Lambda$ of (6), where u_0 is a Fowler solution such that

$$\mathcal{U}(x) = (1 + O(|x|^\alpha))\mathcal{U}_0(x) \quad (4.54)$$

as $x \rightarrow 0$, for some $\alpha > 0$.

Proof. First we observe that (4.53) implies that the origin is a nonremovable singularity. Thus, by Theorem 4.4 we get that $P(\mathcal{U}) < 0$. Consider $v_i(t, \theta)$ the change of variables from Fowler (1.2) of u_i .

Let $\{\tau_k\}$ be a sequence of real numbers such that $\tau_k \rightarrow \infty$. Consider the translated sequence $v_{i,k}(t, \theta) = v_i(t + \tau_k, \theta)$ defined in $(-\tau_k, \infty) \times \mathbb{S}^{n-1}$. By (4.53) we get that

$$c_1 \leq |\mathcal{V}_k(t, \theta)| \leq c_2,$$

where $\mathcal{V}_k = (v_{1,k}, v_{2,k})$. Consequently, by standard elliptic estimates, we get the uniform boundedness of any derivative for $t > 0$. Since $v_{i,k}$ satisfies,

$$\mathcal{L}_{\hat{g}_k}(v_{i,k}) - \sum_{j=1}^2 B_{ij}v_{j,k} + \frac{n(n-2)}{4}|\mathcal{V}_k|^{\frac{4}{n-2}}v_{i,k} = 0,$$

where $\mathcal{L}_{\hat{g}_k}$ and B_{ij} are given by (4.5) and $\hat{g} := dt^2 + d\theta^2 + O(e^{-2t}) \rightarrow dt^2 + d\theta^2$. Standard elliptic estimates imply that there exists a subsequence, also denoted by $v_{i,k}$, which converges in the C_{loc}^2 topology, to a positive solution of

$$\partial_t^2 v_{i,0} + \Delta_{\mathbb{S}^{n-1}} v_{i,0} - \frac{(n-2)^2}{4}v_{i,0} + \frac{n(n-2)}{4}|\mathcal{V}_0|^{\frac{4}{n-2}}v_{i,0} = 0,$$

defined in the whole cylinder. By the characterization result given by Theorem 2.10, such limit is a Fowler-type solution and we know that there exists a Fowler solution v_ε and a vector in the unit sphere with positive coordinates Λ such that $\mathcal{V}_\varepsilon(t) = \Lambda v_\varepsilon(t)$. Hence \mathcal{V}_ε does not depend on θ , and we necessarily have that any angular derivative $\partial_\theta v_{i,k}$ converges uniformly to zero.

Besides, we claim that

$$\begin{aligned} v_{i,k}(t, \theta) &= \bar{v}_{i,k}(t)(1 + o(1)) \\ \nabla v_{i,k}(t, \theta) &= -\bar{v}'_{i,k}(t)(1 + o(1)), \end{aligned}$$

as $t \rightarrow \infty$. In fact, suppose that the first equality above is false. Then there exist $\bar{\varepsilon} > 0$ and sequences $\tau_k \rightarrow \infty$, $\theta_k \rightarrow \theta \in \mathbb{S}^{n-1}$ such that

$$\left| \frac{v_{i,k}(\tau_k, \theta_k)}{\bar{v}_{i,k}(\tau_k)} - 1 \right| \geq \bar{\varepsilon}$$

for some $i \geq 1$. This is a contradiction because, after passing to a subsequence, \mathcal{V}_k converges to a rotationally symmetric Fowler-type solution \mathcal{V}_0 . The second inequality follows from similar arguments.

In the cylindrical setting the Pohozaev integral $P(t, \mathcal{V}) = P(e^{-t}, \mathcal{U})$ becomes

$$P(t, \mathcal{V}) := \int_{t \times \mathbb{S}^{n-1}} \left(\frac{1}{2} |\partial_t \mathcal{V}|^2 - \frac{1}{2} |\nabla_\theta \mathcal{V}|^2 - \frac{(n-2)^2}{8} |\mathcal{V}|^2 + \frac{(n-2)^2}{8} |\mathcal{V}|^{\frac{2n}{n-2}} \right) d\sigma_1.$$

Hence

$$P(\mathcal{V}_\varepsilon) := P(0, \mathcal{V}_\varepsilon) = \lim_{k \rightarrow \infty} P(0, \mathcal{V}_k) = \lim_{k \rightarrow \infty} P(\tau_k, \mathcal{V}) = P(\mathcal{V}). \quad (4.55)$$

So we can conclude that the necksize ε of the limit function is independent of the sequence of numbers τ_k . Therefore, for each sequence $\tau_k \rightarrow \infty$ the correspondent sequence \mathcal{V}_k converges to a function $\mathcal{V}_{\varepsilon, T}(t) = \Lambda v_\varepsilon(t + T)$, with $\Lambda \in \mathbb{S}_+^1$, for some $T \in \mathbb{R}$ which depends on the sequence τ_k .

We will show that there exists $T_0 \in \mathbb{R}$ such that \mathcal{V}_k converges to $\mathcal{V}_{\varepsilon, T_0}$ for any sequence $\tau_k \rightarrow \infty$. The idea is to use a delicate rescaling argument due originally to Leon Simon. In order to do that we will prove several claims using the Jacobi fields studied in subsection 2.5 as a tool.

Let T_ε be the period of \mathcal{V}_ε and $A_\tau = \sup_{t \geq 0} |\partial_\theta \mathcal{V}_\tau|$, where $\mathcal{V}_\tau(t, \theta) = \mathcal{V}(t + \tau, \theta)$. Note that $A_\tau < \infty$, since $|\partial_\theta \mathcal{V}_\tau|$ converges uniformly to zero as $t \rightarrow \infty$.

Claim 1: For every $c > 0$, there exists a positive integer N such that, for any $\tau > 0$, either

(i) $A_\tau \leq ce^{-2\tau}$ or

(ii) A_τ is attained at some point in $I_N \times \mathbb{S}^{n-1}$, where $I_N = [0, NT_\varepsilon]$.

Suppose the Claim is not true. Then there exist a constant $c > 0$ and sequences $\tau_k, s_k \rightarrow \infty, \theta_k \in \mathbb{S}^{n-1}$ such that $|\partial_\theta \mathcal{V}_\tau|(s_k, \theta_k) = A_{\tau_k}$ and $A_{\tau_k} > ce^{-2\tau_k}$ as $k \rightarrow \infty$. Then we can translate back further s_k and define $\tilde{v}_{i,k}(t, \theta) = v_{i,k}(t + s_k, \theta)$. Define $\varphi_{i,k} = A_{\tau_k}^{-1} \partial_\theta \tilde{v}_{i,k}$ and note that $|\varphi_k| \leq 1$, where $\varphi_k = (\varphi_{1,k}, \varphi_{2,k})$. Now, we have

$$\mathcal{L}_{\hat{g}_k}(\tilde{v}_{i,k}) - \sum_j \tilde{B}_{ij} \tilde{v}_{j,k} + \frac{n(n-2)}{4} |\tilde{\mathcal{V}}_k|^{\frac{4}{n-2}} \tilde{v}_{i,k} = 0,$$

where the quantities with tilde are the originals replacing t by $t + \tau_k + s_k$. This implies that

$$L_{cyl}(\tilde{v}_{i,k}) + \frac{n(n-2)}{4} |\tilde{\mathcal{V}}_k|^{\frac{4}{n-2}} \tilde{v}_{i,k} = L_{cyl}(\tilde{v}_{i,k}) - \mathcal{L}_{\hat{g}_k}(\tilde{v}_{i,k}) + \sum_j \tilde{B}_{ij} \tilde{v}_{j,k}.$$

Taking the derivative with respect to θ and multiplying by $A_{\tau_k}^{-1}$, we get

$$\begin{aligned} L_{cyl}(\varphi_{i,k}) + \frac{n(n-2)}{4} |\tilde{\mathcal{V}}_k|^{\frac{4}{n-2}} \varphi_{i,k} + n |\tilde{\mathcal{V}}_k|^{\frac{4}{n-2}-2} \tilde{v}_{i,k} \langle \tilde{\mathcal{V}}_k, \varphi_k \rangle \\ = L_{cyl}(\varphi_{i,k}) - \mathcal{L}_{\hat{g}_k}(\varphi_{i,k}) + \sum_j \tilde{B}_{ij} \varphi_{j,k}. \end{aligned}$$

From (4.6) we have

$$\mathcal{L}_{\hat{g}_k}(\varphi_{i,k}) = \Delta_{\hat{g}_k} \varphi_{i,k} - \frac{(n-2)^2}{4} \varphi_{i,k} - \frac{n-2}{4} e^{-t} \partial_r \log |g_k| \circ \Phi(t, \theta) \varphi_{i,k}.$$

But using the fact $\hat{g} = dt^2 + d\theta^2 + O(e^{-2t})$ and the local expression of the laplacian in this metric, we find that

$$\Delta_{\hat{g}_k} \varphi_{i,k} = \partial_t^2 \varphi_{i,k} + \Delta_{\mathbb{S}^{n-1}} \varphi_{i,k} + O(e^{-2(t+\tau_k+s_k)}).$$

This implies that

$$L_{cyl}(\varphi_{i,k}) + \frac{n(n-2)}{4} |\tilde{\mathcal{V}}_k|^{\frac{4}{n-2}} \varphi_{i,k} + n |\tilde{\mathcal{V}}_k|^{\frac{4}{n-2}-1} \tilde{v}_{i,k} \langle \tilde{\mathcal{V}}_k, \varphi_k \rangle = A_{\tau_k}^{-1} e^{-2(\tau_k+s_k)} O(e^{-2t})$$

where $\varphi_k = (\varphi_{1,k}, \varphi_{2,k})$.

Now we can use elliptic theory to extract a subsequence $\varphi_{i,k}$ which converges in compact subsets to a nontrivial and bounded Jacobi field $\varphi = (\varphi_1, \varphi_2)$ which satisfies the following system

$$L_{cyl}(\varphi_i) + \frac{n(n-2)}{4} v_\varepsilon^{\frac{4}{n-2}} \varphi_i + n\Lambda_i \langle \Lambda, \varphi \rangle v_\varepsilon^{\frac{4}{n-2}} = 0.$$

Since each coordinate function of the limit φ_i has no zero eingencomponent relative to Δ_θ , we get a contradiction because a Jacobi field with such property is necessarily unbounded. This proves the Claim 1.

Now suppose we have a sequence $v_{i,k}(t, \theta)$ converging to $\Lambda_i v_\varepsilon(t + T)$ as $k \rightarrow \infty$. Define

$$w_{i,k}(t, \theta) = v_{i,k}(t, \theta) - \Lambda_i v_\varepsilon(t + T).$$

Set

$$\eta_k = b \max_{I_N} |(w_{1,k}, w_{2,k})|, \quad \bar{\eta}_k = \eta_k + e^{-(2-\delta)\tau_k} \quad \text{and} \quad \varphi_{i,k} = \bar{\eta}_k^{-1} w_{i,k},$$

where $\delta > 0$ is a small number and $b > 0$ is a fixed number to be chosen later. Note that $|(\varphi_{1,k}, \varphi_{2,k})| \leq b^{-1}$ on I_N . Then

$$\mathcal{L}_{\hat{g}_k}(w_{i,k}) + \frac{n(n-2)}{4} \left(|\mathcal{V}_k|^{\frac{4}{n-2}} v_{i,k} - \Lambda_i v_\varepsilon^{\frac{n+2}{n-2}} \right) = E_{i,k} \quad (4.56)$$

where $E_{i,k} = \sum_j B_{ij} v_{j,k} + \Lambda_i (L_{cyl} - \mathcal{L}_{\hat{g}_k}) v_\varepsilon$. First note that by (4.5) we get that $E_{i,k} = O(e^{-2(\tau_k+t)})$ when $t \rightarrow \infty$. Second, observe that

$$|\mathcal{V}_k|^{\frac{4}{n-2}} v_{i,k} - \Lambda_i v_\varepsilon^{\frac{n+2}{n-2}} = |\mathcal{V}_k|^{\frac{4}{n-2}} w_{i,k} + \Lambda_i v_\varepsilon \frac{|\mathcal{V}_k|^{\frac{4}{n-2}} - v_\varepsilon^{\frac{4}{n-2}}}{|\mathcal{V}_k|^2 - v_\varepsilon^2} \sum_j w_{j,k} (v_{j,k} + \Lambda_i v_\varepsilon).$$

Multiplying (4.56) by $\bar{\eta}_k^{-1}$ and taking the limit $k \rightarrow \infty$ we get

$$L_{cyl}(\varphi_i) + \frac{n(n-2)}{4} v_0^{\frac{4}{n-2}} \varphi_i + n\Lambda_i \langle \Lambda, \varphi \rangle v_\varepsilon^{\frac{4}{n-2}} = 0,$$

on the whole cylinder, where $\varphi = (\varphi_1, \varphi_2)$ is a Jacobi field.

Claim 2: The Jacobi field φ is bounded for $t \geq 0$.

To prove this claim we will use the analysis done in subsection 2.5. By the spectral decomposition for the laplacian in the sphere, we know that it is possible to write the Jacobi field as

$$\varphi = a_1 \phi_{\varepsilon,0}^1 + a_2 \phi_{\varepsilon,0}^2 + a_3 \phi_{\varepsilon,0}^3 + a_4 \phi_{\varepsilon,0}^4 + \tilde{\varphi}$$

where $\phi_{\varepsilon,0}^i$ are the linearly independent Jacobi fields corresponding to the eigencomponent independent of θ , and $\tilde{\varphi}$ denotes the projection onto the orthogonal complement. We also know that the functions $\phi_{\varepsilon,0}^1$ and $\phi_{\varepsilon,0}^3$ are bounded and $\phi_{\varepsilon,0}^2$ and $\phi_{\varepsilon,0}^4$ are linearly growing.

Let us show that $\tilde{\varphi}$ is bounded by proving that each $\partial_\theta \tilde{\varphi}_i = \partial_\theta \varphi_i$ is bounded for $t \geq 0$. In fact, the function $\partial_\theta \varphi_i$ is the limit of $\bar{\eta}_k^{-1} \partial_\theta v_{i,k}$, and we can suppose that $\partial_\theta \varphi_i$ is nontrivial, otherwise the result is immediate.

If the first item of Claim 1 happens then

$$\sup_{t \geq 0} (\bar{\eta}_k^{-1} |\partial_\theta v_{i,k}|) \leq \frac{ce^{-2\tau_k}}{\eta_k + e^{-(2-\delta)\tau_k}} \leq C.$$

While if the second item of Claim 1 is true then

$$\sup_{t \geq 0} (\bar{\eta}_k^{-1} |\partial_\theta v_{i,k}|) \leq \sup_{t \geq 0} (\bar{\eta}_k^{-1} |\partial_\theta \mathcal{V}_k|) = \sup_{t \in I_N} (\bar{\eta}_k^{-1} |\partial_\theta \mathcal{V}_k|) \leq C,$$

since the sequence $\bar{\eta}_k^{-1} |\partial_\theta \mathcal{V}_k|$ converges in the C_{loc}^2 topology. Therefore each $\tilde{\varphi}_i$ is bounded for $t \geq 0$, hence exponentially decaying.

To end the proof of the Claim 2 we need to show that $a_2 = a_4 = 0$. To see this note that the convergence $\varphi_{i,k} = \bar{\eta}_k^{-1} w_{i,k} \rightarrow \varphi_i$ implies that

$$\begin{aligned} \mathcal{V}_k &= \Lambda v_{\varepsilon,T} + \bar{\eta}_k \varphi + o(\bar{\eta}_k) \\ &= \Lambda v_{\varepsilon,T} + \bar{\eta}_k (a_1 \phi_{\varepsilon,0}^1 + a_2 \phi_{\varepsilon,0}^2 + a_3 \phi_{\varepsilon,0}^3 + a_4 \phi_{\varepsilon,0}^4 + \tilde{\varphi}) + o(\bar{\eta}_k), \end{aligned}$$

where $v_{\varepsilon,T}(t) = v_\varepsilon(t + T)$. On the other hand by (4.30), (4.55) and the Pohozaev identity, Lemma 4.3, we have that

$$P(0, \mathcal{V}_k) = P(\tau_k, \mathcal{V}) = P(\mathcal{V}) + O(e^{-2\tau_k}) = P(v_{\varepsilon,T}) + O(e^{-2\tau_k}).$$

Since $\lim_{k \rightarrow \infty} (\bar{\eta}_k^{-1} e^{-2\tau_k}) = 0$, we would have a contradiction in case a_2 or a_4 is not zero. Thus each φ is bounded for $t \geq 0$.

Now we will show that there exists some T so that the difference between \mathcal{V} and $\mathcal{V}_{\varepsilon,T} = \Lambda v_{\varepsilon,T}$ goes to zero as $t \rightarrow \infty$.

Since we do not know the correct translation parameter, define $\mathcal{V}_\tau(t, \theta) = \mathcal{V}(t + \tau, \theta)$ and $\mathcal{W}_\tau(t, \theta) = \mathcal{V}_\tau(t, \theta) - \Lambda v_\varepsilon(t)$. Let $C_1 > 0$ be a fixed constant and consider the interval I_N as in the Claim 1. Set also $\eta(\tau) = b \max_{I_N} |\mathcal{W}_\tau|$ and $\bar{\eta}(\tau) = \eta(\tau) + e^{-(2-\delta)\tau}$, where $b > 0$ is a fixed constant to be chosen later. We observe that $\eta(\tau) \rightarrow 0$ as

$\tau \rightarrow \infty$. Let us prove the following claim.

Claim 3: If N , b and τ are sufficiently large and $\bar{\eta}$ is sufficiently small, then there exists s with $|s| \leq C_1 \bar{\eta}(\tau)$ so that $\bar{\eta}(\tau + NT_\varepsilon + s) \leq \frac{1}{2} \bar{\eta}(\tau)$.

Suppose the claim is not true. Then there exists some sequence $\tau_k \rightarrow \infty$ such that $\bar{\eta}(\tau_k) \rightarrow 0$ and for any s satisfying $|s| \leq C_1 \bar{\eta}(\tau_k)$ we have that $\bar{\eta}(\tau_k + NT_\varepsilon + s) > \frac{1}{2} \bar{\eta}(\tau_k)$. Define $\varphi_{i,k} = \bar{\eta}(\tau_k)^{-1} w_{i,\tau_k}$, similarly to the previous claim. We can suppose that $\varphi_{i,k}$ converges in C^∞ on compact sets to a Jacobi field, which by Claim 2 is bounded for $t \geq 0$. So we can write

$$\varphi = a_1 \phi_{\varepsilon,0}^1 + a_3 \phi_{\varepsilon,0}^3 + \tilde{\varphi} \quad (4.57)$$

where $\tilde{\varphi}$ has exponential decay. Note that $|\varphi| \leq b^{-1}$ on I_N , which implies that a_1 and a_3 are uniformly bounded, independently of the sequence τ_k . Moreover, since $\phi_{\varepsilon,0}^1 = v'_\varepsilon \Lambda$ and $\phi_{\varepsilon,0}^3 = v_\varepsilon \bar{\Lambda}$ we have

$$|a_3 v_\varepsilon| \leq |\langle \varphi, \bar{\Lambda} \rangle| + |\langle \tilde{\varphi}, \bar{\Lambda} \rangle| \leq b^{-1} + |\tilde{\varphi}| \quad (4.58)$$

on I_N . We know that $v_\varepsilon \geq \varepsilon$ and $\tilde{\varphi}$ decreases exponentially with a fixed rate, and so we can choose b and N sufficiently large such that $|a_3|$ is sufficiently small.

Set $s_k = -\bar{\eta}(\tau_k) a_1$ whose absolute value is less than $C_1 \bar{\eta}(\tau_k)$ if we choose C_1 sufficiently large. Hence for $t \in [0, 2NT_\varepsilon]$ we have

$$\begin{aligned} \mathcal{W}_{\tau_k+s_k}(t, \theta) &= \mathcal{V}(t + \tau_k - \bar{\eta}(\tau_k) a_1, \theta) - \Lambda v_\varepsilon(t) \\ &= \mathcal{V}_{\tau_k}(t - \bar{\eta}(\tau_k) a_1, \theta) - \Lambda v_\varepsilon(t - \bar{\eta}(\tau_k) a_1) \\ &\quad - \bar{\eta}(\tau_k) a_1 \Lambda \frac{v_\varepsilon(t - \bar{\eta}(\tau_k) a_1) - v_\varepsilon(t)}{-\bar{\eta}(\tau_k) a_1} \\ &= \bar{\eta}(\tau_k) \varphi_k(t - \bar{\eta}(\tau_k) a_1, \theta) - \bar{\eta}(\tau_k) a_1 \phi_{\varepsilon,0}^1 + o(\bar{\eta}(\tau_k)) \\ &= \mathcal{W}_{\tau_k}(t, \theta) - \bar{\eta}(\tau_k) a_1 \phi_{\varepsilon,0}^1 + o(\bar{\eta}(\tau_k)), \end{aligned}$$

where $\varphi_k = (\varphi_{1,k}, \varphi_{2,k})$. Here we used the equality $\mathcal{W}_{\tau_k} = \bar{\eta}(\tau_k) \varphi_k + o(\bar{\eta}(\tau_k))$ and the fact that $\varphi_k(t - \bar{\eta}(\tau_k) a_1, \theta) - \varphi_k(t, \theta)$ goes to zero as $\tau_k \rightarrow \infty$.

Consequently, by (4.57), for $t \in [0, 2NT_\varepsilon]$ we get that

$$\mathcal{W}_{\tau_k+s_k} = \bar{\eta}(\tau_k) \tilde{\varphi} + \bar{\eta}(\tau_k) a_3 \phi_{\varepsilon,0}^3 + o(\bar{\eta}(\tau_k)),$$

which implies

$$\max_{I_N} |\mathcal{W}_{\tau_k+s_k+NT_\varepsilon}| = \max_{[NT_\varepsilon, 2NT_\varepsilon]} |\mathcal{W}_{\tau_k+s_k}| \leq \bar{\eta}(\tau_k) \max_{[NT_\varepsilon, 2NT_\varepsilon]} (|\tilde{\varphi}| + |a_3 v_\varepsilon|) + o(\bar{\eta}(\tau_k)).$$

Since $\tilde{\varphi}$ decreases exponentially with a fixed rate, by (4.58) we can choose N and $b > 0$ sufficiently large in a way that the last equality implies that

$$\max_{I_N} |\mathcal{W}_{\tau_k+s_k+NT_\varepsilon}| \leq \frac{1}{4} \bar{\eta}(\tau_k).$$

On the other hand, note that

$$e^{-(2-\delta)(\tau_k+s_k+NT_\varepsilon)} \leq e^{-(2-\delta)NT_\varepsilon} \bar{\eta}(\tau_k) \leq \frac{1}{4} \bar{\eta}(\tau_k)$$

which implies that $\bar{\eta}(\tau + NT_\varepsilon + s) \leq \frac{1}{2} \bar{\eta}(\tau)$, a contradiction. This ends the proof of the Claim 3.

Once the claim above is proved, using an iterative argument, we are ready to prove that there exists σ such that $w_{i,\sigma} \rightarrow 0$ as $t \rightarrow \infty$ for each coordinate. First choose τ_0 and N sufficiently large satisfying the Claim 3 and such that $C_1 \bar{\eta}(\tau_0) \leq \frac{1}{2} NT_\varepsilon$. Let $s_0 = -\bar{\eta}(\tau_0) a_1$ be chosen as above. Thus we have $|s_0| \leq C_1 \bar{\eta}(\tau_0) \leq \frac{1}{2} NT_\varepsilon$. Define inductively three sequences by

$$\begin{aligned} \sigma_k &= \tau_0 + \sum_{i=0}^{k-1} s_i \\ \tau_k &= \tau_{k-1} + s_{k-1} + NT_\varepsilon = \sigma_k + kNT_\varepsilon \\ s_k &= -\bar{\eta}(\tau_k) a_1. \end{aligned}$$

By the Claim 3 we get by induction $\bar{\eta}(\tau_k) \leq 2^{-k} \bar{\eta}(\tau_0)$ and $|s_k| \leq 2^{-k-1} NT_\varepsilon$. Hence there exists the limit $\sigma = \lim \sigma_k \leq \tau_0 + NT_\varepsilon$ and then $\tau_k \rightarrow \infty$ as $k \rightarrow \infty$.

We claim σ is the correct translation parameter. In fact, choose k such that $t = kNT_\varepsilon + [t]$ with $[t] \in I_N$, and write

$$\begin{aligned} w_{i,\sigma}(t, \theta) &= v_i(t + \sigma, \theta) - \Lambda_i v_\varepsilon(t) \\ &= v_i(t + \sigma, \theta) - v_i(t + \sigma_k, \theta) + v_i(t + \sigma_k, \theta) - \Lambda_i v_\varepsilon(t). \end{aligned}$$

Since $\partial_t v_i$ is uniformly bounded, we have

$$v_i(t + \sigma, \theta) - v_i(t + \sigma_k, \theta) = \partial_t v_i(t_0) \sum_{j=k}^{\infty} s_j = O(2^{-k}),$$

for some t_0 . Besides,

$$v_i(t + \sigma_k, \theta) - \Lambda_i v_\varepsilon(t) = v_i(\tau_k + [t], \theta) - \Lambda_i v_\varepsilon([t]) = w_{i, \tau_k}([t], \theta).$$

Thus,

$$\mathcal{W}_\sigma(t, \theta) = \mathcal{W}_{\tau_k}([t], \theta) + O(2^k).$$

Since $b \max_{I_N} |\mathcal{W}_{\tau_k}| = \eta(\tau_k) \leq \bar{\eta}(\tau_k) \leq 2^{-k} \bar{\eta}(\tau_0)$, it holds $|w_{i, \sigma}(t, \theta)| = O(2^{-k})$ or equivalently, using that $t = kNT_\varepsilon + [t]$, we have

$$|w_{i, \sigma}(t, \theta)| \leq C_1 e^{-\frac{\log 2}{NT_\varepsilon} t}$$

which finishes the proof of the theorem.

■

As a direct consequence of the results proved in this section we have the following corollary.

Corollary 4.11 *Suppose that \mathcal{U} is a solution of the system (1) in the punctured ball $B_1^n(0) \setminus \{0\}$ with the potential A satisfying (H1) and (H2) and $3 \leq n \leq 5$. Then there exist a Fowler-type solution \mathcal{U}_0 from (6) such that*

$$\mathcal{U}(x) = (1 + O(|x|^\alpha)) \mathcal{U}_0(x)$$

as $x \rightarrow 0$, for some $\alpha > 0$.

Bibliography

- [1] ABLOWITZ, M. J., PRINARI, B., AND TRUBATCH, A. D. *Discrete and continuous nonlinear Schrödinger systems*, vol. 302. Cambridge University Press, 2004.
- [2] AKHMEDIEV, N., AND ANKIEWICZ, A. Partially coherent solitons on a finite background. *Physical review letters* 82, 13 (1999), 2661.
- [3] CAFFARELLI, L. A., GIDAS, B., AND SPRUCK, J. Asymptotic symmetry and local behavior of semilinear elliptic equations with critical sobolev growth. *Communications on Pure and Applied Mathematics* 42, 3 (1989), 271–297.
- [4] CHEN, C. C., LIN, C.-S., ET AL. Local behavior of singular positive solutions of semilinear elliptic equations with sobolev exponent. *Duke Mathematical Journal* 78, 2 (1995), 315–334.
- [5] CHEN, Z., AND LIN, C.-S. Removable singularity of positive solutions for a critical elliptic system with isolated singularity. *Mathematische Annalen* 363, 1-2 (2015), 501–523.
- [6] DRUET, O., AND HEBEY, E. Stability for strongly coupled critical elliptic systems in a fully inhomogeneous medium. *Analysis & PDE* 2, 3 (2010), 305–359.
- [7] DRUET, O., HEBEY, E., AND VÉTOIS, J. Bounded stability for strongly coupled critical elliptic systems below the geometric threshold of the conformal laplacian. *Journal of Functional Analysis* 258, 3 (2010), 999–1059.
- [8] ESRY, B., GREENE, C. H., BURKE JR, J. P., AND BOHN, J. L. Hartree-fock theory for double condensates. *Physical Review Letters* 78, 19 (1997), 3594.

- [9] FOWLER, R. Further studies of emden's and similar differential equations. *The Quarterly Journal of Mathematics*, 1 (1931), 259–288.
- [10] GIDAS, B., NI, W.-M., AND NIRENBERG, L. Symmetry and related properties via the maximum principle. *Communications in Mathematical Physics* 68, 3 (1979), 209–243.
- [11] KOREVAAR, N., MAZZEO, R., PACARD, F., AND SCHOEN, R. Refined asymptotics for constant scalar curvature metrics with isolated singularities. *Inventiones mathematicae* 135, 2 (1999), 233–272.
- [12] LI, Y. Y. Prescribing scalar curvature on S^3 , S^4 and related problems. *Journal of functional analysis* 118, 1 (1993), 43–118.
- [13] MARQUES, F. C. Isolated singularities of solutions to the yamabe equation. *Calculus of Variations and Partial Differential Equations* 32, 3 (2008), 349–371.
- [14] MAZZEO, R., AND PACARD, F. Constant scalar curvature metrics with isolated singularities. In *Duke Math. J* (1999), Citeseer.
- [15] MAZZEO, R., POLLACK, D., AND UHLENBECK, K. Moduli spaces of singular yamabe metrics. *Journal of the American Mathematical Society* 9, 2 (1996), 303–344.
- [16] POLLACK, D. Compactness results for complete metrics of constant positive scalar curvature on subdomains of S^n . *Indiana University Mathematics Journal* (1993), 1441–1456.
- [17] REED, M., AND SIMON, B. *IV: Analysis of Operators*, vol. 4. Elsevier, 1978.
- [18] SCHOEN, R. On the number of constant scalar curvature metrics in a conformal class. *Differential geometry* 52 (1991), 311–320.
- [19] SCHOEN, R. M. The existence of weak solutions with prescribed singular behavior for a conformally invariant scalar equation. *Communications on pure and applied mathematics* 41, 3 (1988), 317–392.

- [20] SILVA SANTOS, A. A construction of constant scalar curvature manifolds with Delaunay-type ends. *Ann. Henri Poincaré* 10, 8 (2010), 1487–1535.